

# Copper Metallochaperones

Nigel J. Robinson<sup>1</sup> and Dennis R. Winge<sup>2</sup>

<sup>1</sup>Institute for Cell and Molecular Biosciences, Medical School, Newcastle University, NE2 4HH, United Kingdom; email: N.J.Robinson@ncl.ac.uk

<sup>2</sup>University of Utah Health Sciences Center, Salt Lake City, Utah 84132; email: dennis.winge@hsc.utah.edu

Annu. Rev. Biochem. 2010. 79:537–62

First published online as a Review in Advance on March 5, 2010

The *Annual Review of Biochemistry* is online at [biochem.annualreviews.org](http://biochem.annualreviews.org)

This article's doi:  
10.1146/annurev-biochem-030409-143539

Copyright © 2010 by Annual Reviews.  
All rights reserved

0066-4154/10/0707-0537\$20.00

## Key Words

P<sub>1</sub>-type ATPase, cytochrome oxidase, Cox17, CopZ, Atx1, Ccs1

## Abstract

The current state of knowledge on how copper metallochaperones support the maturation of cuproproteins is reviewed. Copper is needed within mitochondria to supply the Cu<sub>A</sub> and intramembrane Cu<sub>B</sub> sites of cytochrome oxidase, within the *trans*-Golgi network to supply secreted cuproproteins and within the cytosol to supply superoxide dismutase 1 (Sod1). Subpopulations of copper-zinc superoxide dismutase also localize to mitochondria, the secretory system, the nucleus and, in plants, the chloroplast, which also requires copper for plastocyanin. Prokaryotic cuproproteins are found in the cell membrane and in the periplasm of gram-negative bacteria. Cu(I) and Cu(II) form tight complexes with organic molecules and drive redox chemistry, which unrestrained would be destructive. Copper metallochaperones assist copper in reaching vital destinations without inflicting damage or becoming trapped in adventitious binding sites. Copper ions are specifically released from copper metallochaperones upon contact with their cognate cuproproteins and metal transfer is thought to proceed by ligand substitution.

## Contents

INTRODUCTION .....	538	Cox11-Mediated Cu <sub>B</sub> Site Formation .....	550
The Copper Proteome .....	539	Summary of the Key Steps in Copper Metallation of CcO .....	551
Cuproprotein Metallation Reactions .....	539	COPPER METALLOCHAPERONE FOR SUPEROXIDE DISMUTASE .....	551
COPPER METALLOCHAPERONES FOR COPPER TRANSPORTERS .....	541	Role of Ccs1 in Superoxide Dismutase 1 Activation .....	551
Visualization of Adducts with Cytosolic Domains of P <sub>1</sub> -Type ATPases .....	541	Domain Structure of Ccs1 .....	552
The Transfer of Copper .....	542	The Sequence of Events in Ccs1-Mediated Activation of Superoxide Dismutase 1 .....	552
Copper Donation to Membrane Sites .....	544	Ccs1-Independent Activation of Superoxide Dismutase 1 .....	553
Copper Acquisition by Atx1-Like Copper Metallochaperones .....	545	Activation of Superoxide Dismutase 1 in the Mitochondrial Intermembrane Space by Ccs1 ...	553
Handling Copper in a More Oxidizing Periplasm .....	545	Key Features of Ccs1-Mediated Superoxide Dismutase 1 Activation .....	554
COPPER METALLOCHAPERONES FOR CYTOCHROME OXIDASE .....	545	ADDITIONAL TARGETS FOR COPPER METALLOCHAPERONES .....	554
Cox17 .....	545		
Cu-Cox17- and Sco1-Mediated Cu(I) Transfer to Cox2 .....	548		

## INTRODUCTION

As cofactors for enzymes, copper ions are required for processes ranging from oxidative phosphorylation, mobilization of iron, connective tissue cross-linking, pigment formation, neuropeptide amidation, catecholamine synthesis, and antioxidant defense. Copper has additional biological roles that may be distinct from serving as a catalytic moiety in cuproenzymes, for example, in the innate immune response, in the modulation of synaptic transmission, and in angiogenesis. A consequence of these vital functions is that copper deficiency has profound clinical outcomes often associated with neurodegeneration. Dietary intake of copper salts generally exceeds tissue demands, so homeostatic mechanisms exist to modulate uptake and facilitate export through the bile.


Impairment in biliary copper excretion results in liver and brain copper overload in patients with Wilson's disease. Copper's ability to accept and donate single electrons make it an ideal redox cofactor, but copper ions are also complicit in the Fenton reaction and hence capable of driving the generation of deleterious hydroxyl radicals (**Figure 1a**). Cu(II) is located at the top of the Irving-Williams series, and Cu(I) is also highly competitive, and hence, all copper ions have the potential to displace less competitive metals from metalloproteins (**Figure 1b**). For all of these reasons, cellular copper pools must be tightly balanced to sustain a sufficient supply while minimizing toxic effects.

In the 1990s, a class of protein, designated copper metallochaperones, was discovered that functions as intracellular Cu(I) ion shuttles

distributing Cu(I) ions to specific partner proteins, thereby overcoming a high copper chelation capacity of the cytoplasm. In many cells, the abundance of Cu(I)-buffering metallochaperones and glutathione (GSH) molecules results in a cytoplasm depleted of free copper ions. The discoveries of the Atx1 and Ccs1 metallochaperones and the elucidation of their structures were reviewed in the 2001 edition of the *Annual Review of Biochemistry* (1). This review focuses on subsequent advances, including the mechanisms of copper transfer, copper metallochaperone-dependent and -independent maturation of superoxide dismutase and the assembly of the copper sites of cytochrome oxidase.

## The Copper Proteome

Attempts have been made to define the copper proteome (cuproproteome) using genomic sequence data (2, 3). Ten distinct Cu-containing proteins have been identified in prokaryotes, but wide fluctuations exist in the distribution of these proteins within species (2). Analyses of prokaryotic and archaean genomes indicated that nearly 72% of bacteria and 31% of archaean species utilize copper (2). The cuproproteome distribution may actually be wider if novel copper proteins exist. Bioinformatic searches using 39 Cu-binding motif signatures revealed additional putative Cu-binding proteins in Bacteria and Archaea (3). Over 70% of the putative cuproproteins identified in prokaryotes have homologs in Archaea and Eukaryotes. The most frequently utilized cuproprotein is cytochrome *c* oxidase. The second most common cuproprotein is NADH dehydrogenase-2 (2). Secretory cuproenzymes include dopamine- $\beta$ -hydroxylase, tyrosinase, ceruloplasmin, lysyl oxidase, and amine oxidases. Certain bacterial phyla (Thermotogae, Chlorobi, Lactobacillales, and Mollicutes) lack known copper enzymes or copper metallochaperones. However, most of these species contain copper effluxers likely as a defense against the deleterious effect of copper ions in the cytoplasm.



**Figure 1**

- (a) Copper catalyzes production of hydroxyl radicals via the Fenton reaction.  
 (b) Copper has a tendency to form stable complexes relative to other essential divalent metals, and hence it is at the top of the Irving-Williams series.  
 (b) Monovalent copper is also competitive, forming tight protein complexes.


Compartmentalization of the eukaryotic cuproproteome is well established (**Figure 2**). Cuproenzymes are localized within the plasma membrane and the periplasm of gram-negative bacteria. Within plants, copper is required in six compartments, including the cytoplasm, endoplasmic reticulum, mitochondria, chloroplast stroma, thylakoid lumen, and apoplast (4). Copper redistribution between compartments can occur upon transitions to copper deficiency (5).

## Cuproprotein Metallation Reactions

In general, metallation reactions are expected to occur during protein biosynthesis, or shortly thereafter, because many metal cofactors influence protein stability. On cytoplasmic ribosomes, protein biosynthesis and chain elongation are coupled to chaperone-mediated protein folding for many proteins. Nascent polypeptides are protected within the ribosome and can only fold as they emerge from the exit tunnel. Subsequently, molecular chaperones bind nascent polypeptides and actively guide the polypeptides in folding through cycles of binding and release. Whereas cytosolic metalloproteins are likely metallated at this stage, because many cuproenzymes are either compartmentalized or secreted their folding and metallation must occur after translation and translocation. Secretory cuproenzymes pass through the endoplasmic reticulum lumen, where folding reactions commence, and maturation continues during transit through

**Atx1:** the yeast copper metallochaperone for P<sub>1</sub>-type ATPases

**Ccs1:** a copper metallochaperone for Cu-Zn superoxide dismutase



**Figure 2**

Copper is passed (green arrows) to cuproproteins (dark green circles), including metallothioneins Crs5 and Cup1, in for example *Saccharomyces cerevisiae*, by copper metallochaperones (pale green circles, or ovals if membrane associated). Copper permeases (white ovals) import copper into the cell or deliver Cu(I) to compartments, such as the *trans*-Golgi network where secreted cuproproteins (such as Fet3) acquire copper. Fet3 oxidizes ferrous ions (curved brown arrow) to provide substrate for high-affinity iron import by Ftr1. P<sub>1</sub>-type ATPases (such as Ccc2) have one or more soluble N-terminal domains, which engage in regulatory interactions with copper metallochaperones (such as Atx1), but additional Cu(I) donation (hence two arrows) may supply the *trans*-membrane regions for Cu(I) transport. Metallation of the cytochrome *c* oxidase complex (CcO) in mitochondria (OM, outer membrane of mitochondria) involves a pathway via Cox11 for the Cu<sub>B</sub> site of the Cox1 subunit, embedded in the inner membrane (IM), and via Sco1 for the Cu<sub>A</sub> site of Cox2, protruding into the intermembrane space (IMS). Cox17 supplies both pathways. Nuclear encoded mitochondrial proteins, such as Cox17 and Ccs1, are imported across the OM unfolded (pale green line) via the TOM translocase then captured in the IMS, following introduction of disulfide bonds (SS) through the actions of Mia40. A sulfhydryl oxidase Erv1 generates a reactive disulfide on Mia40. A small copper ligand (CuL) supplies Cu(I) (dashed arrows) to the IMS copper metallochaperones.

the Golgi apparatus. Copper metallation occurs within Golgi vesicles.

Proteins discriminate between metal ions imperfectly, at least *in vitro*, yet biology seems to achieve precision in metallation reactions. Metallation is influenced by preferences in coordination geometry and in polarizability of ligand donor atoms. Computational chemistry

studies infer that the specificity of a metal for a set of ligands in a protein cavity depends mainly on the metal's natural abundance in the biological locality, tailored by the properties of the protein and metal (6). For example, two structurally related cyanobacterial proteins, MncA and CucA, are the predominant manganese- and copper-containing proteins in the periplasm

but have similar metal-binding sites (7). The manganoprotein prefers to bind Cu(II), or Cu(I), or Zn(II) rather than Mn(II) in vitro (7). However, precise metallation is achieved by differential compartmentalization during the metallation reactions. The manganoprotein, MncA, is dependent on incorporation of Mn(II) within the cytoplasm, where copper and zinc are highly restricted. Metallated MncA is then secreted into the periplasm as a holoenzyme via the Tat pathway. In contrast, metallation of the copper protein, CuaA, occurs after export of the nascent chain via the Sec system. Another strategy for precision of metallation is through the actions of metallochaperones.

## COPPER METALLOCHAPERONES FOR COPPER TRANSPORTERS

Bacterial *copZ* was discovered in the nucleotide sequences adjacent to a gene encoding a copper-transporting P<sub>1</sub>-type ATPase (8), whereas yeast Atx1 was cloned as a suppressor of oxygen sensitivity in strains lacking superoxide dismutase (9). Studies of these two small proteins (69 residues for CopZ and 73 for Atx1) established the concept of copper metallochaperones. Atx1 and CopZ are similar in amino acid sequence to the N-terminal regions of metal-transporting P<sub>1</sub>-type ATPases, including the one (CopA) from the same operon as CopZ, human Menkes and Wilson ATPases and the P-type ATPase from yeast, Ccc2. All of these proteins have a metal-binding motif MXCXXC (where X is any amino acid) associated with the first loop of a  $\beta\alpha\beta\beta\alpha\beta$  (ferredoxin) structural fold. The single  $\beta\alpha\beta\beta\alpha\beta$ -fold of Atx1 or CopZ is replicated once (CopA), or twice (Ccc2), or six (Menkes and Wilson) times in the cytosolic regions of the ATPases (1). Copper metallochaperones that are variants of this structural theme occur in plants and Archaea. CopZ or Atx1-like copper metallochaperones are not obligatory partners for copper-transporting P<sub>1</sub>-type ATPases since some genomes encode the ATPases but no homologs of these metallochaperones (10).

Yeast mutants deficient in Atx1 are deficient in iron uptake owing to impaired copper supply

to the multicopper oxidase Fet3 (**Figure 2**). Extracellular Fet3 activity is needed to generate ferric substrate for the high-affinity iron importer Ftr1. As noted above, copper metallation of secreted proteins occurs within Golgi vesicles, and this is true of Fet3. The Ccc2 ATPase transports copper into this compartment, and thus Atx1 is phenotypically linked to the activity of the ATPase. Evidence that Atx1 forms associations with the N-terminal region of Ccc2 and that the human homolog Atox1 (sometimes called Hah1) forms associations with the Menkes or Wilson ATPases was obtained from coimmunoprecipitations and Cu-dependent two-hybrid interactions, whereas copper transfer between pairs of recombinant proteins was observed in vitro. A model was proposed in which these copper metallochaperones donate metal ions to the N-terminal regions of the ATPases via a sequence of ligand-exchange reactions. At the time of the 2001 *Annual Review of Biochemistry* article (1), solution structures had been obtained for apo-Atx1, Cu(I)-Atx1, as well as for the apo and Cu(I) forms of one domain of Ccc2 (Ccc2a), plus an X-ray crystal structure was known for the homodimer of Cu(I)-Hah1, among others. The homodimer structure was used to generate models of docked heterodimers (11). Solution structural studies have now examined these anticipated heterodimers.

## Visualization of Adducts with Cytosolic Domains of P<sub>1</sub>-Type ATPases

A solution structure of an Atx1-Cu(I)-Ccc2a heterodimer has been generated (**Figure 3**) (12). The interaction is dependent upon copper, with the first cysteine of each protein (Cys15 of Atx1 and Cys13 of Ccc2a) being essential for complex formation (12). This seems reasonable because Cys15 of Atx1 is the more solvent accessible of its cysteine pair. A collection of lysine residues (residues 24, 28, 62, and 65) confer complementary surface charge to a negative surface of Ccc2a. Conversion of these lysines to glutamate impairs Atx1 activity as


---

**CopZ:** a bacterial protein similar to Atx1 and Atox1

**$\beta\alpha\beta\beta\alpha\beta$ -fold:** ferredoxin-fold metal-binding domains of P<sub>1</sub>-type ATPases and some copper metallochaperones

**Atox1:** the human copper metallochaperone for P<sub>1</sub>-type ATPases

---



**Figure 3**

(a) NMR solution structural model [Protein Data Bank (PDB) code 2GGP] of a heterodimer of the *S. cerevisiae* copper metallochaperone Atx1 and one of two N-terminal cytosolic domains of its cognate P<sub>1</sub>-type copper-transporting ATPase Ccc2. Cysteinyll (yellow) thiol coordinated Cu(I) (sphere) stabilizes a transient complex between similar  $\beta\alpha\beta\beta\alpha\beta$  (ferredoxin)-structural folds (red,  $\alpha$ -helix; blue,  $\beta$ -strand). (b) The interacting faces of Ccc2a and Atx1 have complementary (red, negative; blue, positive) electrostatic potentials.

assessed by loss of yeast two-hybrid interactions with Ccc2a and poor ferrous <sup>55</sup>Fe uptake by yeast strains carrying these Atx1 variants (13). Thus, electrostatics enables encounter complexes to form between the copper metallochaperone and the P<sub>1</sub>-type ATPase (Figure 3). However, the encounter complex must be stabilized by coincident coordination of copper through cysteine residues from both partners if the transient complex is to persist for a sufficient length of time to enable detection by NMR. The complex is also predicted to be transient in vivo as Atx1 is largely found in the cytoplasm rather than associated with Golgi membrane preparations. A possible sequence for the exchange of metal-binding ligands between the two proteins was proposed with Cys13 of Ccc2a first invading a digonally coordinated Cu(I)-Atx1 site after which the second Cys of

Ccc2a replaces the second Cys of Atx1. An expectation is that the heterodimer dissociates upon formation of a bithiolate Cu(I) complex with Ccc2a (Figure 4), although there is only a small change in free energy associated with the transfer of Cu(I) from Atx1 to Ccc2a (14).

### The Transfer of Copper


Copper transfer from Atx1 to Ccc2a in vitro was unaffected by the presence of a 50-fold molar excess of GSH, which was added to chelate any released Cu(I) (14). The off rate from Cu(I)-Atx1 to solvent is thought to be negligible, such that transfer to Ccc2a can only occur by direct protein contact and not via free solution. Presumably, the off rate for copper is enhanced in the heterodimer owing to conformation changes induced in Atx1 by its partner. This predicted switch in the Cu(I)-Atx1 off rate has not been measured, although structural data have suggested what might trigger a change. Loop 5 of Atx1 contains a lysine residue (Lys65), which is conspicuously conserved in many homologs from other species. Lys65 is thought to serve as a counter ion, balancing the negative charge of a metal site in which Cu(I) is bound to two thiolates and shielding the site. In adducts with Ccc2a, this residue is displaced away from the metal-binding site (12). Thus, movement of Lys65 in adducts opens the Atx1 metal site for ligand invasion by Ccc2a. The metal-binding site of apo-Atx1 is disordered, whereas that of apo-Ccc2a is structured (12), perhaps also contributing a small free-energy change to encourage metal release from the former to the latter.

The bacterial copper metallochaperones, CopZ from *Enterococcus hirae* (8, 15) and from *Bacillus subtilis* (16, 17), plus ScAtx1 from *Synechocystis* PCC 6803 (18), have a tendency to multimerize when bound to copper. These copper metallochaperone homodimers are presumed to dissociate to allow heterodimers to form with the ATPases. In the bacterial systems, the electrostatics are reversed relative to yeast Atx1-Cu(I)-Ccc2a, with CopZ and ScAtx1 presenting negative faces to positive ones on the respective P-type ATPase



domains. Bioinformatics initially suggested that the CopA P-type ATPase of *B. subtilis* may import copper; however,  $^{15}\text{N}$  chemical shift differences attributed to CopZ-Cu(I)-CopA showed that the metal-binding region of CopZ adopts an apo-like form and that CopA adopts a Cu(I)-like form in the heterodimer, more consistent with a vector for copper transfer from the copper metallochaperone to the ATPase (17). In support of this prediction,  $\Delta\text{copA}$  *B. subtilis* was subsequently shown to be hypersensitive to elevated copper and deficient in copper export not import (19). Indeed, there is now uncertainty about whether any  $\text{P}_1$ -type ATPases import copper, although this was first proposed for CopA from *E. hirae* and remains the simplest model for cyanobacterial CtaA on the basis of the phenotypes of  $\Delta\text{ctaA}$  strains.

Copper metallochaperones can direct Cu(I) to bona fide destinations but also keep it away from aberrant ones. Zinc-transporting  $\text{P}_1$ -type ATPases, such as ZntA and ZiaA, also possess  $\beta\alpha\beta\alpha\beta$ -folds plus CXXC metal-binding sites, akin to those of copper transporters (20, 21). In ZiaA, from the cyanobacterium *Synechocystis* PCC 6803, the metal site has been shown to have a preference for Cu(I) over Zn(II). However, the N-terminal domain of ZiaA does not form heterodimers with the copper metallochaperone Cu(I)-ScAtx1 (22). Electrostatics repulses encounter complexes between ZiaA and ScAtx1, while complexes between ScAtx1 and related copper transporters CtaA and PacS are not repulsed. This mechanism provides a kinetic barrier to inhibit aberrant association of Cu(I) with ZiaA (21). Whereas most bacteria have no cytoplasmic membranous compartments and no known copper-requiring enzymes in the cytoplasm, cyanobacteria, such as *Synechocystis* PCC 6803, are peculiar in possessing thylakoid membranes housing plastocyanin and a *caa*<sub>3</sub>-type cytochrome oxidase. CtaA, PacS, and ScAtx1 act positively with respect to thylakoid cuproproteins, and crucially, the ATPases are nonredundant (22, 23). PacS has been localized to thylakoid membranes (24), suggesting a model in which CtaA imports copper at plasma membranes while PacS loads



**Figure 4**

Visualization of the Cys-Xaa-Xaa-Cys motifs taken from NMR solution structures (Protein Data Bank codes indicated) of the apo-, Cu(I)-, and heterodimeric forms of Atx1 and Ccc2a to represent the transfer of Cu(I) from Atx1 to Ccc2a via ligand-exchange reactions.

it into thylakoids, analogous to the supply of copper to cuproproteins within the Golgi in eukaryotes. This model, however, relies upon CtaA acting to import copper, which has not been demonstrated for a  $\text{P}_1$ -type ATPase. A credible alternative model is that CtaA transports copper into thylakoids but at a different stage in their biogenesis than when PacS is active. For example, copper must be supplied twice to tyrosinase in mammals, transiently by the Menkes ATPase in Golgi and again by the Menkes ATPase in specialized melanosome compartments derived from the

Golgi (25). Copper binding is necessary to stabilize tyrosinase in the Golgi, but its folding state, coupled with the ionic conditions in the earlier Golgi network, is thought to lead to release of the initial copper atoms.

### Copper Donation to Membrane Sites

Because there is only a small free-energy change associated with metal transfer from copper metallochaperones to the N-terminal regions of P<sub>1</sub>-type ATPase, it has been widely assumed that irreversible membrane translocation driven by the hydrolysis of ATP is what shifts the equilibrium by removing the Cu(I)-ATPase product. However, there is no direct evidence that copper acquired by an N-terminal metal-binding domain of a P<sub>1</sub>-type ATPase is transferred to *trans*-membrane regions for transport. CopZ from *Archaeoglobus fulgidus* interacts with, and can donate metal to, soluble  $\beta\alpha\beta\beta\alpha\beta$ -domains of CopA from the same organism (26). Importantly, Cu(I)-CopZ can also activate ATPase activity of isolated CopA, even when the ATPase is missing its own soluble  $\beta\alpha\beta\beta\alpha\beta$ -domain or when Cu-binding Cys residues have been converted to Ala. In contrast, the purified Cu(I)-bound  $\beta\alpha\beta\beta\alpha\beta$ -domain of CopA, expressed without the rest of the ATPase, does not activate the variant CopA missing the same domain. Under nonturnover conditions (in the absence of Mg<sup>2+</sup>-ATP), irreversible copper transfer can be detected between Cu(I)-CopZ and the membrane metal-binding sites of CopA (26). It is possible that the transfer of metal from copper metallochaperones to the N-terminal regions of P<sub>1</sub>-type ATPases serves regulatory roles and that separate donation events to the metal sites located in the membranous regions might be a common requirement.

The metal-binding domain of CopA interacts with the nucleotide-binding domain of the ATPase, and this interaction is impaired upon Cu(I) binding or nucleotide binding (27, 28). Thus, Cu(I) donation to the N-terminal region could serve to regulate Cu(I) transport by modulating this internal interaction within the

ATPase. It is notable that CopZ from *A. fulgidus* is atypical in structure, having an extra 130-residue N-terminal domain that contains both zinc and a 2Fe2S cluster (29).

The cytosolic domain of the Wilson ATPase containing its ATP-binding site also interacts with the N-terminal  $\beta\alpha\beta\beta\alpha\beta$ -regions *in vitro*, and these interactions are also inhibited by Cu(I) binding (30). Donation of copper to the N-terminal regions may thus expose the ATP-binding site to assist catalytic turnover in response to increased abundance of Cu-bound metallochaperone. Both the Menkes and Wilson ATPases also undergo metal-dependent shifts in their cellular localization when copper becomes elevated, and the N-terminal  $\beta\alpha\beta\beta\alpha\beta$ -regions of the Menkes ATPase are known to influence trafficking (31–33).

The  $V_{\max}$  of CopA from *A. fulgidus* was greater with Cu(I)-CopZ rather than Cu(I) alone (26). The dynamics of copper release from human Atox1 have been investigated by using the Cu(I) chelator bicinchoninic acid (BCA), which forms a Cu(I) BCA<sub>2</sub> complex (34). Transfer progresses via formation of an Atox1-Cu(I)-BCA intermediate before ligand invasion by a second molecule of BCA. Lack of release to GSH presumably implies that the thiol of GSH is unable to invade the Atox1 Cu(I) site in the same manner as BCA. BCA acquires copper more rapidly from Cu(I)-Atox1 than it does from buffer alone because Cu(I)-dithiothreitol complexes dissociate more slowly than the copper metallochaperone. The rate of release to BCA is further enhanced using Atox1 mutant proteins in which alanine residues are substituted for either the shielding lysine or the methionine of the MXCXXC motif. In both cases, rapid release from mutant *Atox1* occurs because the second molecule of BCA more readily displaces Atox1 from the intermediate adduct relative to wild-type Atox1 (34). Copper metallochaperones can thus make copper swiftly accessible to the correct substrates capable of ligand invasion, presumably including membranous metal-binding sites of P<sub>1</sub>-type ATPases.



## Copper Acquisition by Atx1-Like Copper Metallochaperones

Yeast mutants missing any of the diverse types of copper permeases still retain functional copper metallochaperones, implying that there is no single preferred donor (35). Furthermore, because the permeases are so varied, it seems improbable that they can all support specific interactions with copper metallochaperones. The electron crystallographic structure of the Ctr1 copper pore may reveal a cytosolic domain with a candidate docking site for Atx1 (36). Copper transfer between the cytosolic regions of Ctr1 and Atx1 does occur *in vitro*, although the estimated affinities of the two proteins are similar at about  $10^{-19}$  M (37, 38). Constitutive overexpression of competing Cu(I)-binding proteins, such as the metallothioneins Crs5 and Cup1, in the cytosol does not interfere with Cu(I) acquisition by copper metallochaperones. One of the burning questions is how do cells prioritize copper supply? An intriguing notion is that this could be achieved if some, but not all, copper metallochaperones gained access to metal ions released at cuproprotein turnover. Atx1 targets would be subordinate if Atx1 solely obtained copper from *de novo* import, and the primacy of cytochrome oxidase could be sustained provided its copper metallochaperones had preferential access to recycled copper.

## Handling Copper in a More Oxidizing Periplasm

A five-stranded  $\beta$ -barrel protein, CusF, involved in the efflux of surplus copper from the periplasm of gram-negative bacteria, such as *Escherichia coli*, has been added to the catalog of copper metallochaperones. CusF has an unusual copper site, involving a histidine imidazole, two methionine thioethers, plus a  $\pi$ -interaction from the aromatic ring of tryptophan (39–41). The methionine thioether group appears as a Cu ligand in sites located within more oxidizing environments, for example, in copper permeases such as Ctr1 (42). The *E. coli* CusBCA proteins are thought to form a com-

plex that straddles the inner and outer membranes to couple the inner-membrane (IM) proton motive force to copper efflux across the outer membrane. It is predicted that the substrate is acquired in the periplasm rather than the cytoplasm. By using selenomethionine derivatives to distinguish the metal sites, it was possible to follow movement of copper from Cu(I)-SeMet-CusF to apo-CusB by X-ray absorption spectroscopy (43). The proteins have closely matched affinities, leading to a suggestion that metal transfer might be driven by removal of the Cu(I)-CusB product through CusBCA-mediated efflux, analogous to the arguments proposed for the driving force for copper transfer from Atx1 to Ccc2.

## COPPER METALLOCHAPERONES FOR CYTOCHROME OXIDASE

Unlike Atx1 and Ccs1 that donate Cu(I) directly to the target molecule, Cox17-mediated Cu(I) donation to cytochrome *c* oxidase (CcO) subunits involves two accessory factors Cox11 and Sco1, which function in the metallation of the Cu<sub>B</sub> site in Cox1 and Cu<sub>A</sub> site in Cox2, respectively. Both Cox11 and Sco1 are IM-associated proteins with Cu(I)-binding globular domains protruding into the intermembrane space (IMS). Cox17-mediated Cu(I) transfer to Cox11 and Sco1 is the initial event, followed by the subsequent transfer to Cox1 and Cox2, respectively. Cu(I) donation to Cox11 and Sco1 is believed to impart specificity to the Cu(I) transfer to CcO subunits during CcO biogenesis. One difference between the Cu(I) metallation of Cox1 and Cox2 is that the Cu<sub>A</sub> site in Cox2 is localized 8 Å above the IM, whereas the Cu<sub>B</sub> site in Cox1 is buried 13 Å below the membrane surface (44).

### Cox17

Cox17 was cloned from *Saccharomyces cerevisiae* from a CcO deficiency mutant. Cells lacking Cox17 were respiratory deficient and unable to propagate on nonfermentable carbon sources (45). Mitochondria isolated from the mutant

---

**Cox17:** a eukaryotic copper metallochaperone for Sco1 and Cox11


**Cox11:** a copper metallochaperone for the Cu<sub>B</sub> site of cytochrome oxidase

**Sco1:** a copper metallochaperone for the Cu<sub>A</sub> site of cytochrome oxidase

---

cells contained both mitochondrial and nuclear CcO subunits, precluding a role for Cox17 in the expression or import of the subunit polypeptides. The mutant cells lacked the CcO-specific heme *a*, but not heme *b* found in other respiratory complexes or cytochrome *c* (45). Insight into the role of Cox17 in CcO biogenesis was gleaned by the observation that *cox17*Δ cells regained an ability to propagate on glycerol and ethanol medium with the addition of 0.4% copper salts. Because the mutant cells contained a functional Sod1 superoxide dismutase, the role of Cox17 is specific to CcO and/or mitochondria. Mice lacking Cox17 die during early embryogenesis at a stage similar to mice lacking the copper transporter Ctr1 (46, 47). Prior to embryonic death, CcO activity was severely impaired in *COX17*<sup>-/-</sup> mice. Depletion of *COX17* mRNA, using siRNA, in HeLa cells resulted in an induced CcO deficiency (48). Steady-state levels of Cox1 and Cox2 were diminished following the knockdown, but the other respiratory complexes retained the same magnitude of abundance as in wild-type (WT) cells.

Cox17 is a small hydrophilic protein of 63 residues in humans. It contains a conserved twin CX<sub>9</sub>C motif that forms a helical hairpin configuration (Figure 5). This helical hairpin is also referred as a CHCH (coiled-coil<sub>1</sub> helix<sub>1</sub> coiled-coil<sub>2</sub> helix<sub>2</sub>) motif (49). The twin CX<sub>9</sub>C structural motif is found in numerous IMS proteins; several of these, including Cox19, Cox23, and Pet191, are important in CcO biogenesis. The Cox12 structural subunit of CcO projecting into the IMS also contains the twin CX<sub>9</sub>C motif. The cysteines within the twin CX<sub>9</sub>C motif are present in two disulfides, stabilizing the helical hairpin of Cox17 (Figure 5*a*) (50, 51). The redox couple of the double disulfide configuration of Cox17 to the fully reduced state has a midpoint potential of -294 mV, consistent with the dual disulfide molecule being the likely species in vivo, considering that the IMS redox potential is approximately -255 mV unlike the more-reducing potential of the cytoplasm (-296 mV) (52, 53). In support of the disulfide configuration of Cox17 within the IMS is the observation that other Cys-rich




**Figure 5**

Copper metallochaperones for CcO. (a) Disulfide bonds between two pairs of cysteine residues (yellow) on antiparallel  $\alpha$ -helices (red) create a helical hairpin of Cox17 [Protein Data Bank (PDB) code 2RNB]. Cu(I) is coordinated to a pair of cysteine residues. (b) Sco1 is tethered to the inner membrane (IM) (hydrophobic  $\alpha$ -helix not shown) with a thioredoxin fold ( $\beta$ -strand, dark blue; histidine, pale blue) in the intermembrane space (PDB 2GQM). (c) Cox11 is tethered to the IM (hydrophobic  $\alpha$ -helix not shown) with an immunoglobulin-like  $\beta$ -barrel in the IMS (PDB 1SPO).

helical hairpin proteins within the IMS with a twin CX<sub>3</sub>C motif form heterohexameric complexes in which disulfides are essential to stabilize both the structure and function of the complex (55). Although Cox17 likely exists with twin disulfides, Cox17 is functional without either of the two disulfides (54).

Cu(I) binding to Cox17 occurs through two vicinal cysteine residues in a digonal Cu(I)-thiolate complex (**Figures 5a and 6**) (50, 51, 55). The mononuclear Cu(I)-Cox17 has been characterized only after in vitro Cu(I) reconstitution. Recombinant expression of Cox17 in *E. coli* leads to the accumulation of a Cu-Cox17 oligomeric, tetranuclear Cu(I) complex in cells cultured in Cu-supplemented medium (56). Recombinant expression in bacteria without copper supplementation results in the isolation of metal-free protein. This is presumably due to the absence of available cytosolic copper ions in *E. coli* (57). The polycopper protein is in a dimer/tetramer equilibrium, with the Cu(I)-thiolate cluster likely existing at the dimer interface (56). The tetracopper cluster conformer requires multiple cysteine residues to be in the reduced thiolate state. Cox17 purified from the IMS is largely devoid of bound Cu(I) and is monomeric (58). Cu-reconstituted human Cox17 shows a S-Cu-S angle of 130° (55). The bent coordination may permit an exogenous thiolate to provide a third ligand. The Cu(I) thiolate center is charge stabilized by two adjacent Lys residues. Lys20 may mimic the stabilizing Lys65 in the Cu-Atx1 complex. In addition, a hydrophobic patch, formed by ends of the C-terminal helix and the N-terminal segment, orients the Cys thiolates for Cu(I) binding. The Cu(I) center is solvent accessible yet avidly bound with a  $K_d$  value of  $6.4 \times 10^{-15}$  M (59). The Cu(I)-binding affinity must be poised for transfer to target proteins Sco1 and Cox11. Another well-defined digonal Cu(I)-thiolate complex exists in the *E. coli* CueR transcriptional regulator (57). This MerR-based regulator exhibits an essentially linear S-Cu-S bond angle of 176° with short Cu-S bond distances of 2.13 Å. Eight residues separate the two coordinating cysteines. The linearity of the Cu(I)



**Figure 6**

Visualization of metal-binding motifs taken from structures (Protein Data Bank codes indicated) of the apo and Cu(I) forms of Cox17 and Sco1 to represent the transfer of Cu(I) from Cox17 to Sco1 and, potentially, to represent the coincident transfer of two electrons when apo-Sco1 is oxidized.

coordination results in a Cu(I)  $K_d$  near  $10^{-21}$  M. In contrast, the bent coordination of Cu(I) by the vicinal Cys residues in Cox17 reduces the binding affinity perhaps to permit transfer.

The disulfides in Cox17 are introduced during import into the IMS. The IMS import of small cysteine-rich polypeptides with either twin CX<sub>9</sub>C or twin CX<sub>3</sub>C structural motifs is mediated by a disulfide relay system involving the mitochondrial intermembrane space assembly (MIA) machinery consisting of at least

Mia40 and Erv1 (**Figure 2**) (60–64). Transit of these proteins, as unfolded polypeptides through the TOM translocase in the OM, results in transient capture of the imported molecules by Mia40 through an intermolecular disulfide. Mia40 contains two structural disulfide bonds formed by its twin CX<sub>9</sub>C motif as well as a N-terminal labile disulfide. The reactive disulfide lies within a CPC sequence motif and has a reduction potential of –200 mV (65). The nonoptimal stereochemistry of the disulfide with a single residue spacer likely contributes to its instability. The sulfhydryl oxidase Erv1 catalyzes formation of a reactive disulfide in Mia40 that is responsible for transient capture of imported protein with reduced thiolates (60, 61, 66, 67). The second Cys of the CPC motif is the active Cys forming the transient disulfide with Cox17 and other target proteins (65). Protein release from the Mia40 complex is facilitated by disulfide exchange reactions resulting in disulfides in the imported proteins. The oxidative folding of Cox17 and other Mia40 targets traps the proteins within the IMS. It remains unclear whether the Cu(I)-binding Cys23 and Cys24 residues of Cox17 are also initially oxidized. If a disulfide forms with Cys23 and Cys24, reduction would be required for Cu(I) binding. Information is lacking on what redox couples exist within the IMS. GSH reductase and thioredoxin reductase are not known to be present in this compartment.

Cox17 is also present in the yeast cytoplasm, initially encouraging a hypothesis that Cox17 may shuttle Cu(I) to mitochondria (58). However, restricting Cox17 to mitochondria through tethering to the IM retained normal CcO biogenesis (68). Import of Cox17 through the TOM translocase occurs as an unfolded and, presumably, apolypeptide. The source of copper for metallation of Cox1 and Cox2 during CcO biogenesis arises from a mitochondrial matrix copper pool. A significant fraction of mitochondrial copper is localized within the matrix as a soluble, low-molecular-weight ligand complex (CuL), which is conserved from yeast to humans. Information is lacking on whether copper within the CuL complex is

imported as ionic Cu(I) or as a CuL complex. The ligand appears to be present in the apo state within the cytoplasm, so an attractive postulate is that formation of the CuL complex occurs within the cytoplasm and the complex is imported into the mitochondrial matrix through an unknown carrier. Heterologous expression of two Cu(I)-binding proteins in the mitochondrial matrix was shown to titrate out the matrix Cu(I)-ligand complex and induce a respiratory deficiency (69, 70). These effects are reversed by supplementation of cell cultures with CuSO<sub>4</sub>. Attenuation of mitochondrial copper in *Podospora anserina* by a heterologous Cu(I)-binding protein also induces a CcO deficiency (71). In addition, the presence of the matrix-targeted Crs5 in yeast results in diminished Sod1 protein levels within the IMS and impaired activity of an IM-tethered hSod1.

### Cu-Cox17- and Sco1-Mediated Cu(I) Transfer to Cox2

The recipient of Cu(I) from Cu-Cox17 is Sco1 in yeast, and a combination of Sco1 and Sco2 in mammalian cells. Overexpression of Sco1 restored respiratory function in yeast lacking Cox17, thereby linking Cox17 and Sco1 (72). Sco1 participates in the formation of the mixed valent Cu<sub>A</sub> site in Cox2. In the absence of Sco1, yeast are stalled in CcO biogenesis, and the complex is degraded (73, 74). Yeast also possess Sco2, but cells lacking Sco2 are not respiratory deficient. Both human Sco1 and Sco2 proteins are required for CcO biogenesis and have nonoverlapping functions (75). Mutations in either gene results in a fatal encephalomyopathy, although the clinical presentation differs between hSCO1 and hSCO2 patients (76, 77). In addition to the CcO deficiency, patient tissues show a tissue-dependent copper deficiency (78).

Sco1 and Sco2 are tethered to the IM by a single *trans*-membrane helix with a globular domain protruding into the IMS containing a single Cu(I) site (79, 80). Cu(I) is trigonally coordinated by two cysteinyl residues within

a CPDVC sequence motif plus a histidine imidazole (**Figures 5b** and **6**) (80). Mutation of the Cys or His residues abrogates Cu(I) binding, resulting in a nonfunctional CcO (79, 80). The Cu(I) ion is partially shielded by the first Cys and the adjacent two residues (81). The structures of the metal-free human Sco1 and Cu<sub>1</sub>Sco1 complex are similar with the exception of one protruding loop that adopts two different conformations (81) (see the loop marked by an arrow in **Figure 5b**). In the apo conformation, the Cu(I)-binding region is more disordered, and the conformer, frozen in the crystal structure of hSco1, contained the His rotated out of position for planar Cu(I) coordination (82–84). The movement of the protruding loop orients the Cu(I) binding His residue for metal binding. The dynamic properties of this loop suggest it could influence interactions involved in Cu(I) transfer (81).

Human Sco2 shows greater conformational dynamics than hSco1 (85). These proteins have thioredoxin folds, raising questions about whether Sco proteins may function in redox reactions. Recent evidence suggests that hSco2 acts as a thioldisulfide oxidoreductase in CcO biogenesis, whereas hSco1 may be the dominant Cu(I) donor to Cox2 (86). Thioredoxins typically contain a *cis*-Pro in juxtaposition to the redox active cysteinyl residues. The *cis*-Pro precludes metal ion binding (87). The corresponding positions in Sco1 and Sco2 are the Cu(I) ligand His, consistent with their Cu(I)-binding function. If Sco2 is primarily an oxidoreductase, its function remains dependent on the Cys and His residues. Substrate recognition by protein disulfide isomerases, which share the thioredoxin fold (88), may also be dissimilar to partner recognition by copper metallochaperones with the former being dominated by hydrophobic interactions. The reduction potential of the Cys residues in hSco1 and hSco2 are –280 mV and –310 mV, respectively. Considering the reduction potential of the IMS is approximately –255 mV (52), it is predicted that Sco proteins may exist with at least some Cys residues oxidized and therefore not poised for Cu(I) binding. Sco1 and Sco2 in fibroblasts do

have a mixture of oxidized and reduced Cys residues (86).

Direct Cu(I) transfer from Cox17 to Sco1 has been shown *in vitro* and *in vivo*. This likely occurs through a highly transient interaction of the two molecules (59). The interaction presumably imparts specificity to the Cu(I) transfer step. The specificity of the transfer reaction was demonstrated by using mutant proteins (89). Cys57 forms a disulfide with Cys26 in the twin CX<sub>9</sub>C motif of Cox17. Whereas a C<sup>57</sup>S substitution maintains function, a C<sup>57</sup>Y substitution abrogates Cox17 function (45, 54). The nonfunctional C<sup>57</sup>Y Cox17 mutant failed to mediate Cu(I) transfer to Sco1 in both *in vivo* and *in vitro* assays despite binding Cu(I) normally (89). However, the mutant retained the ability to transfer Cu(I) to its second target Cox11. Sco1 P<sup>174</sup>L is an allele of patients with CcO deficiency (90–92). Pro174 is adjacent to the second Cys in the Cu-binding CXXXC motif in Sco1, and this mutant has impaired Cu(I) acquisition from Cox17 (59, 92). A transient interaction persists between P<sup>174</sup>L Sco1 and Cox17, but K<sub>Cu</sub> is weaker, perhaps too weak to allow Cu(I) acquisition from a, now relatively tighter, Cu(I) site in Cox17 (59).

Recently, Cu-Cox17 was shown to effectively transfer both Cu(I) and reducing equivalents to oxidized apo-Sco1, leading to Cu-Sco1 formation and fully oxidized Cox17 (93). The Cu-free conformer of Cox17 failed to reduce oxidized Sco1, suggesting that Cu(I) is necessary for the disulfide exchange mechanism. The ability of Cu-Cox17 to mediate Cu(I) transfer to oxidized Sco1 abrogates a necessary mechanism to maintain reduced thiolates in Sco1. Although Cu-Cox17 can also transfer Cu(I) to Sco2 with reduced thiolates (94), it fails to transfer to oxidized Sco2 (93).

The Cu(I) ions transferred from Cox17 to Sco1 appear to be donated subsequently to Cox2 as an intermediate step in CcO assembly (**Figure 2**). The Sco1-mediated metallation of Cox2 likely occurs after extrusion of the Cox2-soluble domain into the IMS by the Cox18 translocase. Information on the Sco1-mediated metallation of Cox2 in eukaryotes is limited,



although a group of conserved residues on the dynamic protruding loop in Sco1 was shown to be important for function with Cox2 (95). Sco proteins are widely distributed phylogenically. In prokaryotes, Sco1 is often codistributed with CcO and linked to the Cu<sub>A</sub> center in Cox2 (96). However, Sco1 in *Rhodobacter capulatus* is associated with a *cbb*<sub>3</sub> CcO lacking a Cu<sub>A</sub> site (97). The Cu-binding function of this Sco protein, designated SenC, is required for maturation of the *cbb*<sub>3</sub> CcO. One scenario is that SenC is the Cu(I) donor to the Cu<sub>B</sub> center. The *B. subtilis* Sco1 is implicated in formation of the Cu<sub>A</sub> site in CcO. The intriguing aspect of the *Bacillus* Sco1 is its dramatic preference for Cu(II) over Cu(I) (98). The marked stability of the Cu(II)-Sco1 complex may necessitate a conversion step to an intermediate state of Cu(I)-Sco1 compatible with Cu transfer (99).

Cox17 is restricted to eukaryotes. A bioinformatic search for genes associated with Sco1 in bacteria revealed a protein of unknown function with a conserved H(M)X<sub>10</sub>MX<sub>21</sub>HXM sequence motif present in both gram-negative and gram-positive organisms (100). The protein, designated DR1885, has a single-candidate *trans*-membrane helix followed by a globular domain folded into a Greek key  $\beta$ -barrel conformation capable of binding a single Cu(I) ion with a Met<sub>3</sub>His ligand set. The protein is analogous to other bacterial periplasmic Cu-binding proteins that use Met-rich motifs to coordinate Cu(I). The surface-exposed Cu(I) site makes it accessible for Cu(I) transfer. A homologous protein in *Thermus*, designated PCu<sub>A</sub>C for the periplasmic Cu<sub>A</sub> chaperone, was shown to bind Cu(I) with a subpicomolar affinity (101). Cu-PCu<sub>A</sub>C was able to transfer Cu(I) in vitro to the reduced, soluble Cu<sub>A</sub> domain of CcO subunit II of *Thermus* without the assistance of Sco1 (101). Successive Cu(I) transfer steps assembled the binuclear Cu<sub>A</sub> center in subunit II. No transfer occurred when the Cys residues in subunit II were oxidized, unless reduced apo-Sco1 was present, leading to the oxidation of Sco1. Thus, *Thermus* Sco1 functions as an oxidoreductase, rather than a Cu(I) donor, to form the Cu<sub>A</sub> center in CcO. The di-

rect Cu(I) transfer from Cu-PCu<sub>A</sub>C to subunit II may involve ligand-exchange steps with the two Cys residues in the Cu<sub>A</sub> center. Cu(I) coordination by a Met<sub>3</sub>His protein is preferred over a Cys-based donor like Cox17 in the more oxidizing periplasm. The DR1885/PCu<sub>A</sub>C family of proteins may functionally replace Cox17 in prokaryotes as Cu(I) donors to Cu<sub>A</sub> centers that face the periplasmic space. A clear prediction is that bacterial cells lacking DR1885/PCu<sub>A</sub>C may be impaired in CcO biogenesis.

The implications of Sco1 having a redox role rather than a Cu(I) transfer function in *Thermus* are unclear. Maintaining Sco1 in the reduced state competent for a redox function would require reducing equivalents in the periplasm. In yeast, Sco1 has a Cu(I) transfer function, and it remains to be demonstrated whether it possesses a redox function. In humans, Sco2 is implicated in a redox function (86).

### Cox11-Mediated Cu<sub>B</sub> Site Formation

Copper metallation of the Cu<sub>B</sub> site in Cox1 requires Cox11 (Figure 2). *S. cerevisiae* lacking Cox11 have impaired CcO activity and less Cox1 (102). CcO isolated from *Rhodobacter sphaeroides* *cox11*Δ cells lacked Cu<sub>B</sub> but contained both hemes, and the environment of the heme in the Cu<sub>B</sub>-heme *a*<sub>3</sub> site was altered (103). The Cu<sub>A</sub> site in *R. sphaeroides* CcO was unaffected in cells lacking Cox11. Thus, the absence of Cox11 appears to specifically preclude Cu<sub>B</sub> site formation.

Cox11 is tethered to the IM by a single *trans*-membrane helix with a C-terminal domain protruding into the IMS (104, 105). The Cu(I)-binding cysteinyl residues lie within this C-terminal domain. The structure of the globular domain of the Cox11 homolog from *Simorbiobium meliloti* adopts an immunoglobulin-like  $\beta$ -barrel fold (Figure 5c) (106). Removal of the *trans*-membrane domain of yeast Cox11 yields a soluble protein that dimerizes upon Cu(I) binding (104). The Cu(I) sites in each monomer are closely juxtaposed as the dimeric complexes can form a binuclear Cu(I) thiolate cluster at the dimer interface. Mutations of two Cys residues

within a CXC sequence motif abrogate Cu(I) binding and formation of active CcO. A third functionally important Cys is spatially removed from the CXC motif, yet when mutated, this Cys results in substoichiometric Cu(I) binding.

Cu<sub>B</sub> site formation occurs in Cox1 prior to addition of the Cox2 or Cox3 subunits. The Cu<sub>B</sub>-heme *a*<sub>3</sub> binuclear site forms in an early assembly intermediate of Cox1 that is stabilized by the Shy1 assembly factor. Cox11 forms a transient interaction with Shy1, which may be important to coordinate Cu<sub>B</sub> site formation. This proposition is supported by the observation that accumulation of the nonfunctional CcO complexes in *Rhodobacter* and *Paracoccus surf1*Δ (Shy1 homolog) cells is compromised in both Cu<sub>B</sub> and heme *a*<sub>3</sub> (107, 108).

The membrane-embedded Cu<sub>B</sub> site in Cox1 consists of three His ligands lying within a 12-helical bundle. In the absence of Cox2, a partially occluded channel forms from the IMS side of Cox1, where the heme *a* and Cu(I) moieties enter. The putative physical transfer from Cu-Cox11 to the buried Cu<sub>B</sub> site in Cox1 may involve the third Cys residue that was removed from the CXC motif. This Cys is predicted to be juxtaposed to the IM and may mediate the Cu(I) transfer. However, it is not clear whether it would mediate Cu(I) transfer through the channel or laterally through an accessible port of the helical bundle. Cox11 appears to transiently occlude the Cox1 channel, as mutant alleles lacking the Cu-binding CXC residues confer resistance to the pro-oxidant heme *a*<sub>3</sub>:Cox1 assembly intermediate (109).

### Summary of the Key Steps in Copper Metallation of CcO

Copper-transfer reactions are initiated within the IMS by Cu-Cox17, which acquires Cu(I) from a small molecule (**Figure 2**). Cox17 transfers Cu(I) ions to IM-associated Sco1 and Cox11. Cu-Cox11 is the donor to the Cu<sub>B</sub> site in Cox1, and this site is formed in a Cox1 assembly intermediate that may contain only the Cox5a and Cox6 peripheral CcO subunits. The assembly intermediate is fully inserted within

the IM and is associated with CcO assembly factors, including Shy1. Upon hemylation of the two heme *a* centers and formation of Cu<sub>B</sub>, Cox1 is poised for the addition of Cox2. It is not clear whether the binuclear Cu<sub>A</sub> site in Cox2 is formed in an isolated Cox2 or as Cox2 is associated with metallated Cox1. Sco1 mediates Cu<sub>A</sub> site formation, and this reaction may occur as a ternary complex of Cu-Cox17, Sco1, and Cox2. Once the redox copper and heme cofactors are added to Cox1 and Cox2, the remaining subunits are added for final CcO maturation.

### COPPER METALLOCHAPERONE FOR SUPEROXIDE DISMUTASE

The copper metallochaperone for Cu,Zn-superoxide dismutase is Ccs1, a 27-kDa three-domain polypeptide (**Figure 2**) (110, 111). Two of the domains, domains 1 and 3, bind Cu(I), whereas the central domain is a key domain for interaction with Sod1 (112, 113). The physiological significance of Ccs1 has been defined in multiple species. Yeast lacking Ccs1 is devoid of Sod1 enzymatic activity unless cells are cultured with high levels of copper (114, 115). *Drosophila melanogaster* deficient in Ccs1 lacks Sod1 activity and shows exquisite sensitivity to redox cycling of paraquat (116). Mice with a targeted disruption of *CCS* have a markedly attenuated level of Sod1 activity and exhibit no apparent radio-copper incorporation into Sod1 (117). Thus, a strong correlation exists that Ccs1 is required for Sod1 activation.

### Role of Ccs1 in Superoxide Dismutase 1 Activation

Cu-Ccs1 induces the activation of monomeric reduced Sod1, facilitating both Cu(I) transfer and disulfide bond formation (118). Active Sod1 requires binding of both Zn(II) and Cu and formation of an intrasubunit disulfide bond between Cys57 and Cys146 (118, 119). Sod1 is unusual in possessing such an essential disulfide bond in the reducing cytoplasm. The reduction potential of the Sod1 disulfide is

−230 mV, whereas the potential in the cytoplasm is approximately −290 mV. The stability of the Sod1 disulfide may arise from the low solvent accessibility of Cys146 in the disulfide pair as well as from the stabilization imparted by Sod1 dimerization. The copper center in Sod1 is sunken within an active site channel that superoxide anions enter for dismutation (120, 121). In the absence of the metal cofactors and upon reduction of the disulfide, the Sod1 dimer is destabilized and exists as an inactive monomer (118, 122). Metal binding and/or disulfide bond formation stabilizes the  $\beta$ -barrel and loop conformations within a dimeric enzyme.

### Domain Structure of Ccs1

The N-terminal domain 1 of Ccs1 is an Atx1-like  $\beta\alpha\beta\beta\alpha\beta$ -fold capable of binding a single Cu(I) ion (123). Domain 1 of yeast Ccs1 is required for Sod1 activation in Cu-limited cells, but this domain is not essential under normal growth conditions (114). In the absence of domain 1, no radiocopper incorporation into mouse Sod1 was observed in cultured mouse fibroblasts, but copper salts restored limited active Sod1 (124). *Drosophila* and *Anopheles gambiae* Ccs1 proteins lack domain 1, yet these proteins activate Sod1 efficiently (116). Human mutant Ccs1 with Cys substitutions in the MXCXXC motif and Cu(I) bound to domain 3 retain the in vitro activation function of Sod1 (123).

Domain 2 adopts an eight-stranded  $\beta$ -barrel structure analogous to Sod1 (125, 126). The dimeric domain 2 in human Ccs1 contains a Zn(II) site similar to Sod1 but lacks the ligand set for Cu coordination. In contrast, yeast Ccs1 domain 2 lacks both the Zn and Cu centers. Domain 2 is important for docking with Sod1 during the activation reaction. Mutations that compromise Ccs1 dimerization also abrogate both Ccs1 and Sod1 interaction and Ccs1-mediated Sod1 activation (111, 124).

Domain 3 is a short C-terminal segment containing a critical CXC motif that is unstructured in the yeast apo-Ccs1 protein (125), but domain 3 is ordered in the co-structure of yeast

Ccs1 and a mutant Sod1 that stabilizes the intermolecular complex (113). Ccs1 lacking the domain 3 CXC motif fails to transfer Cu(I) or to induce disulfide formation in Sod1 (114, 123, 124). The binding of Ccs1 to Sod1 involves both domains 2 and 3, and binding is retained if the Cys residues are mutated (111). Whereas the domain 3 CXC motif may suggest digonal Cu(I)-thiolate coordination, EXAFS analyses of the Cu(I) complex clearly showed a polycopper cluster. The cluster formed with only domain 3 Cys residues is consistent with a binuclear Cu(I)-thiolate center bridging two Ccs1 molecules (123). Polycopper cluster formation is lost when the domain 3 Cys residues are mutated.

### The Sequence of Events in Ccs1-Mediated Activation of Superoxide Dismutase 1

The Ccs1-Sod1 complex is transient and becomes detectable with reduced Sod1 lacking a Cu site or in mutant Sod1 lacking one Cys of the disulfide pair (112, 118, 127). In the absence of bound Cu in Sod1, complex formation between Ccs1 and Sod1 is dependent on the disulfide being reduced. A loop in Sod1 containing Cys57, one of the disulfide-forming residues, is more flexible when the disulfide bond is reduced (128). Ccs1 is postulated to recognize a partially folded nascent conformation of Sod1 through mobile loop elements (127). Ccs1 interaction with Sod1 is abrogated when the loops become ordered in the copper-bound, disulfide-bonded fold. The Ccs1 interaction with Sod1 is independent of Zn(II) binding to Sod1. Because Zn binding to Sod1 stabilizes a folded conformation (129), Ccs1 must be competent to interact with a partially folded Sod1 conformer, but this structure has reduced thiolates and an empty copper site. The activation process involves copper insertion and disulfide bond formation.

Cu site formation appears to be concurrent with disulfide bond formation. Sod1 mutants blocked in Cu site formation remain largely in the reduced state in yeast containing Ccs1

(127). Likewise, *in vitro* studies revealed that reduced Zn-Sod1 protein is inefficiently converted to the oxidized state aerobically in the absence of Ccs1 (118). Sod1 oxidation requires the intact CXC motif in Ccs1 (130). The addition of Cu-Ccs1 to Zn-Sod1 fails to induce disulfide bond formation anaerobically, but this process is facile with the addition of air (118). A putative intermediate in the oxidation reaction is an intermolecular disulfide between Cys57 of Sod1 and Cys229 of Ccs1 (113). Formation of this intermolecular disulfide may resolve to the C57-C146 disulfide by a disulfide exchange reaction.

### Ccs1-Independent Activation of Superoxide Dismutase 1

Although Ccs1 has a key role in Sod1 activation, it is now clear that a secondary mechanism exists for Sod1 activation independent of Ccs1 in multicellular eukaryotes. The *Caenorhabditis elegans* genome completely lacks Ccs1 (131). Mice lacking Ccs1 have 10% to 30% WT Sod1 activity in various tissues (132). Likewise, Ccs1-null flies show residual Sod1 activity such that the phenotype of the null flies is less severe compared to flies lacking Sod1 (116). Human Sod1 is capable of being partially activated independently of Ccs1 when expressed in yeast or flies (116, 133). This Ccs1-independent pathway of Sod1 activation does not require Atx1 or Cox17 (133).

Yeast Sod1 differs from human and fly Sod1 in depending on Ccs1 for activation (115, 131). Two C-terminal Pro residues in yeast Sod1 preclude Ccs1-independent activation. Replacement of the yeast Pro residues restores Ccs1-independent activation, and insertion of Pro residues at the corresponding positions in human Sod1 abrogates Ccs1-independent activation (133). The Ccs1-independent pathway requires GSH, although the role of GSH in Cu insertion and/or disulfide bond formation is unclear. Because Sod1 proteins lacking the C-terminal Pro residues are Cu activated in *ccs1Δ* yeast cells (116, 131), there must be some bioavailable Cu(I) complexes in the yeast cy-

tosol. The identity of the bioavailable Cu(I) complex is a major unresolved question. Cu metallation of Sod1 by such complexes is less efficient than by Cu-Ccs1, but once it occurs, disulfide bond formation also proceeds. The facile oxidation of worm Sod1 in the absence of Ccs1 implies a secondary pathway (131). Disulfide bond formation is typically a facilitated process, but sulfhydryl oxidases are not known in the cytoplasm.

### Activation of Superoxide Dismutase 1 in the Mitochondrial Intermembrane Space by Ccs1

Ccs1 is key to activation of Sod1 within the mitochondrial IMS, where 1% to 5% of Sod1 resides (**Figure 2**). The presence of Sod1 within the IMS is largely dependent on Ccs1 (134). In the absence of Ccs1, only minimal levels of Sod1 are apparent in the IMS (131). The limiting factor for the import of Ccs1 into the IMS is Mia40 (135–137). Overexpression of Mia40 in yeast or mammalian cells enhances IMS levels of Ccs1, and depletion of Mia40 has the opposite effect (136, 137). Import and capture of Ccs1 in the IMS are likely achieved through disulfide bonding to Mia40, followed by transfer of the disulfide to Ccs1 either in domain 1 or 3 (137). Two factors question whether oxidative folding of Ccs1 is the major mechanism of its retention within the IMS. First, it is unclear whether disulfide bonding within the CXXC in domain 1 or CXC in domain 3 would be sufficient to trap Ccs1 within the IMS as the cross-links would not stabilize dispersed segments of the Ccs1 polypeptide to sterically block diffusion back out through the TOM translocase. Second, the stability of those candidate disulfides may be low because of unsatisfactory spacing between Cys residues (138). This also supports a transient disulfide introduced in domain 1 of Ccs1 upon IMS import. However, retention of Ccs1 in the IMS must involve general folding of domains 1 and 2. Because the Zn site in domain 2 is not conserved in all Ccs1 orthologs, metal binding is not an obvious trigger for IMS folding.

If transient disulfides form in domain 1 of Ccs1, it is unlikely this disulfide is transferred initially to imported Sod1 as Cu insertion must precede disulfide bond formation in Sod1. It is conceivable that Cu(I) insertion into Sod1 is mediated by domain 3 of Ccs1 and that the disulfide is subsequently transferred from the Ccs1 domain 1. Sod1 is not an apparent substrate of the Mia40/Erv1 system (137), so disulfide bond formation in Sod1 must occur through a Ccs1-dependent pathway as in the cytoplasm. Human Sod1 contains two additional Cys residues (C6 and C111), and neither of these is involved in a disulfide bond, yet retention of human Sod1 in the mitochondria of cultured cells is dependent on both (136). Cys111 is near the dimer interface, and the distance between the two Cys side chains is about 10 Å. Cys111 was reported to be part of a nonnative Cu(II)-binding site, leading to speculation that it may mediate Cu insertion into the active site (139).

### Key Features of Ccs1-Mediated Superoxide Dismutase 1 Activation

Ccs1 binds Cu(I) in either domains 1 or 3. Domain 1 binding must necessitate transfer to the domain 3 CXC motif for subsequent transfer to Sod1. Cu-charged Ccs1 interacts with newly synthesized Sod1 polypeptides prior to completion of its native fold. The transient interaction is mediated by domains 2 and 3 of Ccs1. Zn(II) binding to Sod1 can occur either prior to Ccs1 complex formation or concomitantly. Cu(I) transfer occurs from Cu(I) associated in the domain 3 CXC motif. The only characterized Cu(I)-bound state in domain 3 is the binuclear Cu(I) thiolate cluster bridging a Ccs1 dimer. A second candidate Cu(I) donor is a mononuclear digonal Cu(I) complex, but this state has not been reported to date. Cu(I) transfer to Sod1 is accompanied by disulfide bond formation in an oxygen- or superoxide-dependent manner. Major unresolved questions are the mechanism by which the disulfide bond is formed and the sequence of Cu(I)-ligand exchange. The major questions in the Ccs1-

independent pathway of Sod1 activation include the identity of the bioavailable copper pool(s) used in Sod1 metallation and the mechanism of disulfide bond formation.

### ADDITIONAL TARGETS FOR COPPER METALLOCHAPERONES

Atx1 is implicated in copper supply to amine oxidase 1 (Cao1) in *Schizosaccharomyces pombe*, which has been localized (as a green fluorescent protein fusion) to the cytosol rather than the Golgi or some other internal membranous compartment (140). Of the copper metallochaperones, only disruption of Atx1 caused a loss of enzymatic activity, and a yeast two-hybrid interaction has been detected between Atx1 and Cao1. *Arabidopsis thaliana* has a second Atx1, CCH, with an extra C-terminal domain that is inhibitory to its interactions with the P<sub>1</sub>-type ATPases HMA5 (4, 141–143). CCH is expressed in the vicinity of xylem and phloem elements, and one hypothesis is that the C-terminal region enables movement through plasmodesmata, suggesting a role for CCH in systemic Cu trafficking.

Mutants of *E. birae* missing CopZ show low expression of the *cop* operon, including *copA* and *copB*, plus hypersensitivity to copper resulting from insufficient copper-exporting P<sub>1</sub>-type ATPase activity (8). If CopZ solely donated Cu(I) ions to a copper exporter, then intracellular copper levels would be elevated in  $\Delta copZ$ , and copper-responsive expression of the *cop* operon would be predicted to become higher. An alternate target for CopZ has been suggested. Transcriptional regulation of the *cop* operon is mediated by the DNA-binding repressor CopY. Purified CopY contains one Zn(II) ion, which can be displaced by two copper ions to form a Cu(I)-thiolate cluster (144, 145). Binding of CopY to nucleotide sequences from the *cop* operator promoter region is impaired by the replacement of zinc with copper. The second  $\beta\alpha\beta\beta\alpha\beta$ -domain of the Menkes ATPases expressed in isolation is incapable of donating Cu(I) to CopY, although a variant in which four residues are



replaced with lysines does donate (145). This supports the idea of direct protein contact mediated by complementary electrostatic surfaces on CopZ and CopY. As with analogous Cu(I) transfer experiments, the metal might not attain a similar equilibrium by exchanging via free solution owing to its kinetic properties, notably the copper metallochaperone off rate for Cu(I), that is unique to the heterodimer interface.

Atox1 can affect the transcriptional control of cyclin D1 and mouse embryonic fibroblast proliferation (146). A yeast two-hybrid interaction also occurs between Ccs1 and the  $\beta$ -secretase (BACE1) that cleaves the amyloid precursor protein and generates the A $\beta$  peptides that aggregate in Alzheimer's disease se-

nile plaques. Ccs1 and BACE1 can be coimmunoprecipitated in extracts from rat brains (147). A short cytosolic tail of BACE1 contains three Cys residues that bind Cu(I), and this same short region associates with domain 1 of Ccs1. Fluorescent fusions of the two proteins track together along axons of neuronal cells, heading toward the synapse, implying that Ccs1 forms a stable association with secretory vesicles containing BACE1. Perhaps the intermolecular interaction between domain 1 of Ccs1 and BACE1's tail is structurally similar to transient intramolecular interactions with domain 3 of Ccs1. Some detected copper metallochaperone interactions might not be physiological, but aberrations linked to grave pathologies demand consideration.

## SUMMARY POINTS

1. The switch between cuprous and cupric ions is exploited in oxidoreductases but can also catalyze deleterious redox chemistry, for example via the Fenton reaction.
2. Cu(II) is at the top of the Irving-Williams series, and Cu(I) also has a tendency to form tight thiol complexes; hence copper is liable to outcompete other metals in metalloproteins.
3. The cytosol contains numerous potential copper-binding sites, and the estimated available copper concentration is negligible, probably subfemtomolar.
4. Copper is passed to cuproproteins by ligand-exchange reactions from copper metallochaperones to avoid copper becoming kinetically trapped, or engaging in deleterious interactions while en route with docking between a metallochaperone and its partner, presumed to encourage the release of copper from the metallochaperone.
5. Solution structural studies have visualized transient complexes between N-terminal domains of P<sub>1</sub>-type ATPases and Atox1-type copper metallochaperones to reveal changes predicted to aid Cu(I) release and to reveal interacting surfaces with complementary electrostatics.
6. Interactions between Atox1-type copper metallochaperones and N termini of P<sub>1</sub>-type ATPase are regulatory, and Cu(I) can be separately donated to ATPase membrane sites.
7. Cox17, Sco1, and Cox11 in the mitochondrial IMS form two copper supply pathways for Cu<sub>A</sub> and Cu<sub>B</sub>, sites of cytochrome oxidase.
8. Ccs1 introduces an essential disulfide bond into Sod1 and supplies Cu(I) to Sod1 both in the cytosol and in the mitochondrial IMS, although several Sod1 species can acquire Cu(I) and the critical disulfide without Ccs1 provided GSH is present.

## FUTURE ISSUES

1. Studies over the coming years should aim to further visualize heterodimer complexes between Ccs1 and Sod1, the mitochondrial copper metallochaperones for cytochrome oxidase and their partners, along with CopZ and membranous regions of P<sub>1</sub>-type ATPases.
2. The effect that docking between a metallochaperone and its partner has on K<sub>OFF</sub> for Cu(I) from the metallochaperone needs to be quantified, perhaps by stopped-flow methods.
3. In addition to determining the sequence of Cu(I)-ligand exchange, proton and electron migration that prepares recipient Cys residues for Cu(I) awaits a full description, as does the coupling of Cu(I) transfer from Ccs1 to Sod1 with disulfide bond formation in Sod1.
4. Additional twin CX<sub>9</sub>C proteins within the IMS contribute to CcO biogenesis, including Cox19, Cox23, and Pet191, but their actions remain to be defined.
5. The sources of copper for each metallochaperone remain to be unequivocally defined, the low-molecular-weight mitochondrial matrix copper ligand needs to be structurally characterized, and mechanisms that prioritize different cuproprotein destinations are yet to be discovered.

## DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

## ACKNOWLEDGMENTS

The authors are supported by the Biotechnology and Biological Sciences Research Council (BB/E001688/1) to N.J.R. and grant ES 03817 from the National Institutes of Environmental Health Sciences, NIH to D.R.W.

## LITERATURE CITED

1. Huffman DL, O'Halloran TV. 2001. Function, structure, and mechanism of intracellular copper trafficking proteins. *Annu. Rev. Biochem.* 70:677–701
2. Ridge PG, Zhang Y, Gladyshev VN. 2008. Comparative genomic analyses of copper transporters and cuproproteomes reveal evolutionary dynamics of copper utilization and its link to oxygen. *PLoS ONE* 3:e1378
3. Andreini C, Banci L, Bertini I, Rosato A. 2008. Occurrence of copper proteins through the three domains of life: a bioinformatic approach. *J. Proteome Res.* 7:209–16
4. Burkhead JL, Reynolds KA, Abdel-Ghany SE, CoHu CM, Pilon M. 2009. Copper homeostasis. *New Phytol.* 182:799–816
5. Merchant SS, Allen MD, Kropat J, Moseley JL, Long JC, et al. 2006. Between a rock and a hard place: trace element nutrition in *Chlamydomonas*. *Biochim. Biophys. Acta.* 1763:578–94
6. Dudev T, Lim C. 2008. Metal binding affinity and selectivity in metalloproteins: insights from computational studies. *Annu. Rev. Biophys.* 37:97–116
7. Tottey S, Waldron KJ, Firbank SJ, Reale B, Bessant C, et al. 2008. Protein-folding location can regulate manganese-binding versus copper- or zinc-binding. *Nature* 455:1138–42
8. Odermatt A, Solioz M. 1995. Two *trans*-acting metalloregulatory proteins controlling expression of the copper-ATPases of *Enterococcus hirae*. *J. Biol. Chem.* 270:4349–54

9. Lin SJ, Culotta VC. 1995. The ATX1 gene of *Saccharomyces cerevisiae* encodes a small metal homeostasis factor that protects cells against reactive oxygen toxicity. *Proc. Natl. Acad. Sci. USA* 92:3784–88
10. Zhang Y, Gladyshev VN. 2009. Comparative genomics of trace elements: emerging dynamic view of trace element utilization and function. *Chem. Rev.* 109:4828–61
11. Wernimont AK, Huffman DL, Lamb AL, O'Halloran TV, Rosenzweig AC. 2000. Structural basis for copper transfer by the metallochaperone for the Menkes/Wilson disease proteins. *Nat. Struct. Biol.* 7:766–71
12. Banci L, Bertini I, Cantini F, Felli IC, Gonnelli L, et al. 2006. The Atx1-Ccc2 complex is a metal-mediated protein-protein interaction. *Nat. Chem. Biol.* 2:367–68
13. Portnoy ME, Rosenzweig AC, Rae T, Huffman DL, O'Halloran TV, Culotta VC. 1999. Structure-function analyses of the ATX1 metallochaperone. *J. Biol. Chem.* 274:15041–45
14. Huffman DL, O'Halloran TV. 2000. Energetics of copper trafficking between the Atx1 metallochaperone and the intracellular copper transporter, Ccc2. *J. Biol. Chem.* 275:18611–14
15. Wimmer R, Herrmann T, Solioz M, Wuthrich K. 1999. NMR structure and metal interactions of the CopZ copper chaperone. *J. Biol. Chem.* 274:2597–603
16. Zhou L, Singleton C, Le Brun NE. 2008. High Cu(I) and low proton affinities of the CXXC motif of *Bacillus subtilis* CopZ. *Biochem. J.* 413:459–65
17. Banci L, Bertini I, Ciofi-Baffoni S, Del Conte R, Gonnelli L. 2003. Understanding copper trafficking in bacteria: interaction between the copper transport protein CopZ and the N-terminal domain of the copper ATPase CopA from *Bacillus subtilis*. *Biochemistry* 42:1939–49
18. Banci L, Bertini I, Ciofi-Baffoni S, Kandias NG, Robinson NJ, et al. 2006. The delivery of copper for thylakoid import observed by NMR. *Proc. Natl. Acad. Sci. USA* 103:8320–25
19. Radford DS, Kihlken MA, Borrelly GP, Harwood CR, Le Brun NE, Cavet JS. 2003. CopZ from *Bacillus subtilis* interacts in vivo with a copper exporting CPx-type ATPase CopA. *FEMS Microbiol. Lett.* 220:105–12
20. Banci L, Bertini I, Ciofi-Baffoni S, Finney LA, Outten CE, O'Halloran TV. 2002. A new zinc-protein coordination site in intracellular metal trafficking: solution structure of the apo and Zn(II) forms of ZntA (46–118). *J. Mol. Biol.* 323:883–97
21. Banci L, Bertini I, Ciofi-Baffoni S, Poggi L, Vanarotti M, et al. 2010. NMR structural analysis of the soluble domain of ZiaA-ATPase and the basis of selective interactions with copper metallochaperone Atx1. *J. Biol. Inorg. Chem.* 15:87–98
22. Tottey S, Rondet SA, Borrelly GP, Robinson PJ, Rich PR, Robinson NJ. 2002. A copper metallochaperone for photosynthesis and respiration reveals metal-specific targets, interaction with an importer, and alternative sites for copper acquisition. *J. Biol. Chem.* 277:5490–97
23. Tottey S, Rich PR, Rondet SAM, Robinson NJ. 2001. Two Menkes-type ATPases supply copper for photosynthesis in *Synechocystis* PCC 6803. *J. Biol. Chem.* 276:19999–20004
24. Kanamaru K, Kashiwagi S, Mizuno T. 1994. A copper-transporting P-type ATPase found in the thylakoid membrane of the cyanobacterium *Synechococcus* species PCC7942. *Mol. Microbiol.* 13:369–77
25. Setty SR, Tenza D, Sviderskaya EV, Bennett DC, Raposo G, Marks MS. 2008. Cell-specific ATP7A transport sustains copper-dependent tyrosinase activity in melanosomes. *Nature* 454:1142–46
26. González-Guerrero M, Arguello JM. 2008. Mechanism of Cu<sup>+</sup>-transporting ATPases: soluble Cu<sup>+</sup> chaperones directly transfer Cu<sup>+</sup> to transmembrane transport sites. *Proc. Natl. Acad. Sci. USA* 105:5992–97
27. Wu CC, Rice WJ, Stokes DL. 2008. Structure of a copper pump suggests a regulatory role for its metal-binding domain. *Structure* 16:976–85
28. Gonzalez-Guerrero M, Hong D, Arguello JM. 2009. Chaperone-mediated Cu<sup>+</sup> delivery to Cu<sup>+</sup> transport ATPases: requirement of nucleotide binding. *J. Biol. Chem.* 284:20804–11
29. Sazinsky MH, LeMoine B, Orofino M, Davydov R, Bencze KZ, et al. 2007. Characterization and structure of a Zn<sup>2+</sup> and [2Fe-2S]-containing copper chaperone from *Archaeoglobus fulgidus*. *J. Biol. Chem.* 282:25950–59
30. Tsvikovskii R, MacArthur BC, Lutsenko S. 2001. The Lys1010-Lys1325 fragment of the Wilson's disease protein binds nucleotides and interacts with the N-terminal domain of this protein in a copper-dependent manner. *J. Biol. Chem.* 276:2234–42

31. Petris MJ, Mercer JFB, Culvenor JG, Lockhart P, Gleeson PA, Camakaris J. 1996. Ligand-regulated transport of the Menkes copper P-type ATPase efflux pump from the Golgi apparatus to the plasma membrane: a novel mechanism of regulated trafficking. *EMBO J.* 15:6084–95
32. Voskoboinik I, Strausak D, Greenough M, Brooks H, Petris M, et al. 1999. Functional analysis of the N-terminal CXXC metal-binding motifs in the human Menkes copper-transporting P-type ATPase expressed in cultured mammalian cells. *J. Biol. Chem.* 274:22008–12
33. Guo Y, Nyasae L, Braiterman LT, Hubbard AL. 2005. NH<sub>2</sub>-terminal signals in ATP7B Cu-ATPase mediate its Cu-dependent anterograde traffic in polarized hepatic cells. *Am. J. Physiol. Gastrointest. Liver Physiol.* 289:G904–16
34. Hussain F, Olson JS, Wittung-Stafshede P. 2008. Conserved residues modulate copper release in human copper chaperone Atox1. *Proc. Natl. Acad. Sci. USA* 105:11158–63
35. Portnoy ME, Schmidt PJ, Rogers RS, Culotta VC. 2001. Metal transporters that contribute copper to metallochaperones in *Saccharomyces cerevisiae*. *Mol. Genet. Genomics* 265:873–82
36. De Feo CJ, Aller SG, Siluvai GS, Blackburn NJ, Unger VM. 2009. Three-dimensional structure of the human copper transporter hCTR1. *Proc. Natl. Acad. Sci. USA* 106:4237–42
37. Xiao Z, Wedd AG. 2002. A C-terminal domain of the membrane copper pump Ctr1 exchanges copper(I) with the copper chaperone Atx1. *Chem. Commun.* 6:588–89
38. Xiao Z, Loughlin F, George GN, Howlett GJ, Wedd AG. 2004. C-terminal domain of the membrane copper transporter Ctr1 from *Saccharomyces cerevisiae* binds four Cu(I) ions as a cuprous-thiolate polynuclear cluster: sub-femtomolar Cu(I) affinity of three proteins involved in copper trafficking. *J. Am. Chem. Soc.* 126:3081–90
39. Loftin IR, Franke S, Blackburn NJ, McEvoy MM. 2007. Unusual Cu(I)/Ag(I) coordination of *Escherichia coli* CusF as revealed by atomic resolution crystallography and X-ray absorption spectroscopy. *Protein Sci.* 16:2287–93
40. Loftin IR, Franke S, Roberts SA, Weichsel A, Heroux A, et al. 2005. A novel copper-binding fold for the periplasmic copper resistance protein CusF. *Biochemistry* 44:10533–40
41. Xue Y, Davis AV, Balakrishnan G, Stasser JP, Staehlin BM, et al. 2008. Cu(I) recognition via cation- $\pi$  and methionine interactions in CusF. *Nat. Chem. Biol.* 4:107–9
42. Davis AV, O'Halloran TV. 2008. A place for thioether chemistry in cellular copper ion recognition and trafficking. *Nat. Chem. Biol.* 4:148–51
43. Bagai I, Rensing C, Blackburn NJ, McEvoy MM. 2008. Direct metal transfer between periplasmic proteins identifies a bacterial copper chaperone. *Biochemistry* 47:11408–14
44. Tsukihara T, Aoyama H, Yamashita E, Tomizaki T, Yamaguchi H, et al. 1995. Structures of metal sites of oxidized bovine heart cytochrome c oxidase at 2.8 Å. *Science* 269:1069–74
45. Glerum DM, Shtanko A, Tzagoloff A. 1996. Characterization of COX17, a yeast gene involved in copper metabolism and assembly of cytochrome oxidase. *J. Biol. Chem.* 271:14504–9
46. Takahashi Y, Kako K, Kashiwabara S-I, Takehara A, Inada Y, et al. 2002. Mammalian copper chaperone Cox17 has an essential role in activation of cytochrome c oxidase and embryonic development. *Mol. Cell. Biol.* 22:7614–21
47. Lee J, Prohaska JR, Thiele DJ. 2001. Essential role for mammalian copper transporter Ctr1 in copper homeostasis and embryonic development. *Proc. Natl. Acad. Sci. USA* 98:6842–47
48. Oswald C, Krause-Buchholz U, Rödel G. 2009. Knockdown of human COX17 affects assembly and supramolecular organization of cytochrome c oxidase. *J. Mol. Biol.* 389:470–79
49. Banci L, Bertini I, Ciofi-Baffoni S, Tokatlidis K. 2009. The coiled coil-helix-coiled coil-helix proteins may be redox proteins. *FEBS Lett.* 583:1699–702
50. Abajian C, Yatsunyk LA, Ramirez BE, Rosenzweig AC. 2004. Yeast Cox17 solution structure and copper(I) binding. *J. Biol. Chem.* 279:53584–92
51. Arnesano F, Balatri E, Banci L, Bertini I, Winge DR. 2005. Folding studies of Cox17 reveal an important interplay of cysteine oxidase and copper binding. *Structure* 13:713–22
52. Hu J, Dong L, Outten CE. 2008. The redox environment in the mitochondrial intermembrane space is maintained separately from the cytosol and matrix. *J. Biol. Chem.* 283:29126–34
53. Palumaa P, Zovo K. 2008. Modulation of redox switches of copper chaperone Cox17 by Zn(II) ions, determined by new ESI MS based approach. *Antioxid. Redox Signal.* 11:985–95

54. Heaton D, Nittis T, Srinivasan C, Winge DR. 2000. Mutational analysis of the mitochondrial copper metallochaperone Cox17. *J. Biol. Chem.* 275:37582–87
55. Banci L, Bertini I, Ciofi-Baffoni S, Janicka A, Martinelli M, et al. 2008. A structural-dynamical characterization of human Cox17. *J. Biol. Chem.* 283:7912–20
56. Heaton DN, George GN, Garrison G, Winge DR. 2001. The mitochondrial copper metallochaperone Cox17 exists as an oligomeric polycopper complex. *Biochem.* 40:743–51
57. Changela A, Chen K, Xue Y, Holschen J, Outten CE, et al. 2003. Molecular basis of metal-ion selectivity and zeptomolar sensitivity by CueR. *Science* 301:1383–87
58. Beers J, Glerum DM, Tzagoloff A. 1997. Purification, characterization, and localization of yeast Cox17p, a mitochondrial copper shuttle. *J. Biol. Chem.* 272:33191–96
59. Banci L, Bertini I, Ciofi-Baffoni S, Leontari I, Martinelli M, et al. 2007. Human Sco1 functional studies and pathological implications of the P174L mutant. *Proc. Natl. Acad. Sci. USA* 104:15–20
60. Rissler M, Wiedemann N, Pfannschmidt S, Gabriel K, Guiard B, et al. 2005. The essential mitochondrial protein Erv1 cooperates with Mia40 in biogenesis of intermembrane proteins. *J. Mol. Biol.* 353:485–92
61. Chacinska A, Pfannschmidt S, Wiedemann N, Kozjak V, Sanjuan Szklarz LK, et al. 2004. Essential role of Mia40 in import and assembly of mitochondrial intermembrane space proteins. *EMBO J.* 23:3735–46
62. Allen S, Balabanidou V, Sideris DP, Lisowsky T, Tokatidis K. 2005. Erv1 mediates the Mia40-dependent protein import pathway and provides a functional link to the respiratory chain by shuttling electrons to cytochrome *c*. *J. Mol. Biol.* 353:937–44
63. Tokatlidis K. 2005. A disulfide relay system in mitochondria. *Cell* 121:965–67
64. Terziyska N, Grumbt B, Bien M, Neupert W, Herrmann JM, Hell K. 2007. The sulfhydryl oxidase Erv1 is a substrate of the Mia40-dependent protein translocation pathway. *FEBS Lett.* 581:1098–102
65. Banci L, Bertini I, Cefaro C, Ciofi-Baffoni S, Gallo A, et al. 2009. MIA40 is an oxidoreductase that catalyzes oxidative protein folding in mitochondria. *Nat. Struct. Mol. Biol.* 16:198–206
66. Mesecke N, Terziyska N, Kozany C, Baumann F, Neupert W, et al. 2005. A disulfide relay system in the intermembrane space of mitochondria that mediates protein import. *Cell* 121:1059–69
67. Terziyska N, Lutz T, Kozany C, Mokranjac D, Mesecke N, et al. 2005. Mia40, a novel factor for protein import into the intermembrane space of mitochondria is able to bind metal ions. *FEBS Lett.* 579:179–84
68. Maxfield AB, Heaton DN, Winge DR. 2004. Cox17 is functional when tethered to the mitochondrial inner membrane. *J. Biol. Chem.* 279:5072–80
69. Cobine PA, Ojeda LD, Rigby KM, Winge DR. 2004. Yeast contain a non-proteinaceous pool of copper in the mitochondrial matrix. *J. Biol. Chem.* 279:14447–55
70. Cobine PA, Pierrel F, Bestwick ML, Winge DR. 2006. Mitochondrial matrix copper complex used in metallation of cytochrome oxidase and superoxide dismutase. *J. Biol. Chem.* 281:36552–59
71. Scheckhuber CQ, Grief J, Boilan E, Luce K, Debacq-Chainiaux F, et al. 2009. Age-related cellular copper dynamics in the fungal ageing model *Podospora anserina* and in ageing human fibroblasts. *PLoS ONE* 4:e4919
72. Glerum DM, Shtanko A, Tzagoloff A. 1996. *SCO1* and *SCO2* act as high copy suppressors of a mitochondrial copper recruitment defect in *Saccharomyces cerevisiae*. *J. Biol. Chem.* 271:20531–35
73. Schulze M, Rödel G. 1988. *SCO1*, a yeast nuclear gene essential for accumulation of mitochondrial cytochrome *c* oxidase subunit II. *Mol. Gen. Genet.* 211:492–98
74. Krummeck G, Rödel G. 1990. Yeast *SCO1* protein is required for a post-translational step in the accumulation of mitochondrial cytochrome *c* oxidase subunits I and II. *Curr. Genet.* 18:13–15
75. Leary SC, Kaufman BA, Pellechia G, Gguercin G-H, Mattman A, et al. 2004. Human *SCO1* and *SCO2* have independent, cooperative functions in copper delivery to cytochrome *c* oxidase. *Hum. Mol. Genet.* 13:1839–48
76. Shoubridge EA. 2001. Cytochrome *c* oxidase deficiency. *Am. J. Med. Genet.* 106:46–52
77. Stiburek L, Vesela K, Hansikova H, Hulkova H, Zeman J. 2009. Loss of function of Sco1 and its interaction with cytochrome *c* oxidase. *Am. J. Physiol. Cell Physiol.* 296:C1218–26
78. Leary SC, Cobine PA, Kaufman BA, Guercin GH, Mattman A, et al. 2007. The human cytochrome *c* oxidase assembly factors *SCO1* and *SCO2* have regulatory roles in the maintenance of cellular copper homeostasis. *Cell Metab.* 5:9–20



79. Rentzsch N, Krummeck-Weiß G, Hofer A, Bartuschka A, Ostermann K, Rödel G. 1999. Mitochondrial copper metabolism in yeast: mutational analysis of Sco1p involved in the biogenesis of cytochrome *c* oxidase. *Curr. Genet.* 35:103–8
80. Nittis T, George GN, Winge DR. 2001. Yeast Sco1, a protein essential for cytochrome *c* oxidase function is a Cu(I)-binding protein. *J. Biol. Chem.* 276:42520–26
81. Banci L, Bertini I, Calderone V, Ciofi-Baffoni S, Mangani S, et al. 2006. A hint for the function of human Sco1 from different structures. *Proc. Natl. Acad. Sci. USA* 103:8595–600
82. Williams JC, Sue C, Banting GS, Yang H, Glerum DM, et al. 2005. Crystal structure of human SCO1: implications for redox signaling by a mitochondrial cytochrome *c* oxidase “assembly” protein. *J. Biol. Chem.* 280:15202–11
83. Abajian C, Rosenzweig AC. 2006. Crystal structure of yeast Sco1. *J. Biol. Inorg. Chem.* 11:459–66
84. Banci L, Bertini I, Calderone V, Ciofi-Baffoni S, Mangani S, et al. 2006. A hint for the function of human Sco1 from different structures. *Proc. Natl. Acad. Sci. USA* 103:8595–600
85. Banci L, Bertini I, Ciofi-Baffoni S, Gerothanassis IP, Leontari I, et al. 2007. A structural characterization of human SCO2. *Structure* 15:1132–40
86. Leary SC, Sasarman F, Nishimura T, Shoubridge EA. 2009. Human SCO2 is required for the synthesis of CO II and as a thiol-disulphide oxidoreductase for SCO1. *Hum. Mol. Genet.* 18:2230–40
87. Su D, Berndt C, Fomenko DE, Holmgren A, Gladyshev VN. 2007. Conserved cis-proline precludes metal binding by the active site thiolates in members of the thioredoxin family of proteins. *Biochemistry* 46:6903–10
88. Riemer J, Bulleid N, Herrmann JM. 2009. Disulfide formation in the ER and mitochondria: two solutions to a common process. *Science* 324:1284–87
89. Horng YC, Cobine PA, Maxfield AB, Carr HS, Winge DR. 2004. Specific copper transfer from the Cox17 metallochaperone to both Sco1 and Cox11 in the assembly of yeast cytochrome *c* oxidase. *J. Biol. Chem.* 279:35334–40
90. Valnot I, Osmond S, Gigarel N, Mehaye B, Amiel J, et al. 2000. Mutations of the *SCO1* gene in mitochondrial cytochrome *c* oxidase deficiency with neonatal-onset hepatic failure and encephalopathy. *Am. J. Hum. Genet.* 67:1104–9
91. Paret C, Lode A, Krause-Buchholz U, Rödel G. 2000. The P(174)L mutation in the human *bSCO1* gene affects the assembly of cytochrome *c* oxidase. *Biochem. Biophys. Res. Commun.* 279:341–47
92. Cobine PA, Pierrel F, Leary SC, Sasarman F, Horng YC, et al. 2006. The P174L mutation in human Sco1 severely compromises Cox17-dependent metallation but does not impair copper binding. *J. Biol. Chem.* 281:12270–76
93. Banci L, Bertini I, Ciofi-Baffoni S, Hadjiloi T, Martinelli M, Palumaa P. 2008. Mitochondrial copper(I) transfer from Cox17 to Sco1 is coupled to electron transfer. *Proc. Natl. Acad. Sci. USA* 105:6803–8
94. Horng YC, Leary SC, Cobine PA, Young FB, George GN, et al. 2005. Human Sco1 and Sco2 function as copper-binding proteins. *J. Biol. Chem.* 280:34113–22
95. Rigby K, Cobine PA, Khalimonchuk O, Winge DR. 2008. Mapping the functional interaction of Sco1 and Cox2 in cytochrome oxidase biogenesis. *J. Biol. Chem.* 283:15015–22
96. Banci L, Bertini I, Cavallaro G, Rosato A. 2007. The functions of Sco proteins from genome-based analysis. *J. Proteome Res.* 6:1568–79
97. Swem DL, Swem LR, Setterdahl A, Bauer CE. 2005. Involvement of SenC in assembly of cytochrome *c* oxidase in *Rhodobacter capsulatus*. *J. Bacteriol.* 187:8081–87
98. Davidson DE, Hill BC. 2009. Stability of oxidized, reduced and copper bound forms of *Bacillus subtilis* Sco. *Biochim. Biophys. Acta.* 1794:275–81
99. Cawthorn TR, Poulsen BE, Davidson DE, Andrews D, Hill BC. 2009. Probing the kinetics and thermodynamics of copper(II) binding to *Bacillus subtilis* Sco, a protein involved in the assembly of the Cu(A) center of cytochrome *c* oxidase. *Biochemistry* 48:4448–54
100. Banci L, Bertini I, Ciofi-Baffoni S, Katsari E, Katsaros N, et al. 2005. A copper(I) protein possibly involved in the assembly of Cu<sub>A</sub> center of bacterial cytochrome *c* oxidase. *Proc. Natl. Acad. Sci. USA* 102:3994–99
101. Abriata LA, Banci L, Bertini I, Ciofi-Baffoni S, Gkazonis P, et al. 2008. Mechanism of Cu(A) assembly. *Nat. Chem. Biol.* 4:599–601

102. Tzagoloff A, Capitanio N, Nobrega MP, Gatti D. 1990. Cytochrome oxidase assembly in yeast requires the product of COX11, a homolog of the *P. denitrificans* protein encoded by ORF3. *EMBO J.* 9:2759–64
103. Hiser L, Di Valentin M, Hamer AG, Hosler JP. 2000. Cox11 is required for stable formation of the Cu<sub>B</sub> and magnesium centers of cytochrome *c* oxidase. *J. Biol. Chem.* 275:619–23
104. Carr HS, George GN, Winge DR. 2002. Yeast Cox11, a protein essential for cytochrome *c* oxidase assembly, is a Cu(I) binding protein. *J. Biol. Chem.* 277:31237–42
105. Carr HS, Maxfield AB, Horng Y-C, Winge DR. 2005. Functional analysis of the domains of Cox11. *J. Biol. Chem.* 280:22664–69
106. Banci L, Bertini I, Cantini F, Ciofi-Baffoni S, Gonnelli L, Mangani S. 2004. Solution structure of Cox11: a novel type of  $\beta$ -immunoglobulin-like fold involved in Cu<sub>B</sub> site formation of cytochrome *c* oxidase. *J. Biol. Chem.* 279:34833–39
107. Smith D, Gray J, Mitchell L, Antholine WE, Hosler JP. 2005. Assembly of cytochrome *c* oxidase in the absence of the assembly protein Surf1p leads to loss of the active site heme. *J. Biol. Chem.* 280:17652–56
108. Bundschuh FA, Hoffmeier K, Ludwig B. 2008. Two variants of the assembly factor Surf1 target specific terminal oxidases in *Paracoccus denitrificans*. *Biochim. Biophys. Acta* 1777:1336–43
109. Khalimonchuk O, Bird A, Winge DR. 2007. Evidence for a pro-oxidant intermediate in the assembly of cytochrome oxidase. *J. Biol. Chem.* 282:17442–49
110. Culotta VC, Klomp LWJ, Strain J, Casareno RLB, Krems B, Gitlin JD. 1997. The copper chaperone for superoxide dismutase. *J. Biol. Chem.* 272:23469–72
111. Schmidt PJ, Kunst C, Culotta VC. 2000. Copper activation of superoxide dismutase 1 (SOD1) in vivo. *J. Biol. Chem.* 275:33771–76
112. Lamb AL, Torres AS, O'Halloran TV, Rosenzweig AC. 2000. Heterodimer formation between superoxide dismutase and its copper chaperone. *Biochemistry* 39:14720–27
113. Lamb AL, Torres AS, O'Halloran TV, Rosenzweig AC. 2001. Heterodimeric structure of superoxide dismutase in complex with its metallochaperone. *Nat. Struct. Biol.* 8:751–55
114. Schmidt PJ, Rae TD, Pufahl RA, Hamma T, Strain J, et al. 1999. Multiple protein domains contribute to the action of the copper chaperone for superoxide dismutase. *J. Biol. Chem.* 274:23719–25
115. Rae RD, Schmidt PJ, Pufahl RA, Culotta VC, O'Halloran TV. 1999. Undetectable intracellular free copper: the requirement of a copper chaperone for superoxide dismutase. *Science* 284:805–7
116. Kirby K, Jensen LT, Binnington J, Hilliker AJ, Ulloa J, et al. 2008. Instability of superoxide dismutase 1 of *Drosophila* in mutants deficient for its cognate copper chaperone. *J. Biol. Chem.* 283:35393–401
117. Wong PC, Waggoner D, Subramaniam JR, Tessarollo L, Bartnikas TB, et al. 2000. Copper chaperone for superoxide dismutase is essential to activate mammalian Cu/Zn superoxide dismutase. *Proc. Natl. Acad. Sci. USA* 97:2886–91
118. Furukawa Y, Torres AS, O'Halloran TV. 2004. Oxygen-induced maturation of SOD1: a key role for disulfide formation by the copper chaperone CCS. *EMBO J.* 23:2872–81
119. Fridovich I. 1997. Superoxide anion radical (O<sub>2</sub><sup>-</sup>), superoxide dismutases and related matters. *J. Biol. Chem.* 272:18515–17
120. Tainer JA, Getzoff ED, Beem KM, Richardson JS, Richardson DC. 1982. Determination and analysis of the 2 A-structure of copper, zinc superoxide dismutase. *J. Mol. Biol.* 160:181–217
121. Tainer JA, Getzoff ED, Richardson JS, Richardson DC. 1983. Structure and mechanism of copper, zinc superoxide dismutase. *Nature* 306:284–87
122. Forman HJ, Fridovich I. 1973. On the stability of bovine superoxide dismutase. The effects of metals. *J. Biol. Chem.* 248:2645–49
123. Stasser JP, Siluvai GS, Barry AN, Blackburn NJ. 2007. A multinuclear copper(I) cluster forms the dimerization interface in copper-loaded human copper chaperone for superoxide dismutase. *Biochemistry* 46:11845–56
124. Caruano-Yzermans AL, Bartnikas TB, Gitlin JD. 2006. Mechanisms of the copper-dependent turnover of the copper chaperone for superoxide dismutase. *J. Biol. Chem.* 281:13581–87
125. Lamb AL, Wernimont AK, Pufahl RA, O'Halloran TV, Rosenzweig AC. 1999. Crystal structure of the copper chaperone for superoxide dismutase. *Nat. Struct. Biol.* 6:724–29
126. Lamb AL, Wernimont AK, Pufahl RA, O'Halloran TV, Rosenzweig AC. 2000. Crystal structure of the second domain of the human copper chaperone for superoxide dismutase. *Biochemistry* 39:1589–95

127. Winkler DD, Schuermann JP, Cao X, Holloway SP, Borchelt DR, et al. 2009. Structural and biophysical properties of the pathogenic SOD1 variant H46R/H48Q. *Biochemistry* 48:3436–47
128. Banci L, Bertini I, Cantini F, D'Onofrio M, Viezzoli MS. 2002. Structure and dynamics of copper-free SOD: the protein before binding copper. *Protein Sci.* 11:2479–92
129. Potter SZ, Zhu H, Shaw BF, Rodriguez JA, Doucette PA, et al. 2007. Binding of a single zinc ion to one subunit of copper-zinc superoxide dismutase apoprotein substantially influences the structure and stability of the entire homodimeric protein. *J. Am. Chem. Soc.* 129:4575–83
130. Proescher JB, Son M, Elliott JL, Culotta VC. 2008. Biological effects of CCS in the absence of SOD1 enzyme activation: implications for disease in a mouse model for ALS. *Hum. Mol. Genet.* 17:1728–37
131. Jensen LT, Culotta VC. 2005. Activation of CuZn superoxide dismutases from *Caenorhabditis elegans* does not require the copper chaperone CCS. *J. Biol. Chem.* 280:41373–79
132. Subramaniam JR, Lyons WE, Liu J, Bartnikas TB, Rothstein J, et al. 2002. Mutant SOD1 causes motor neuron disease independent of copper chaperone-mediated copper loading. *Nat. Neurosci.* 5:301–7
133. Carroll MC, Girouard JB, Ulloa JL, Subramaniam JR, Wong PC, et al. 2004. Mechanisms for activating Cu- and Zn-containing superoxide dismutase in the absence of the CCS Cu chaperone. *Proc. Natl. Acad. Sci. USA* 101:5964–69
134. Field LS, Furukawa Y, O'Halloran TV, Culotta VC. 2003. Factors controlling the uptake of yeast copper/zinc superoxide dismutase into mitochondria. *J. Biol. Chem.* 278:28052–59
135. Son M, Puttaparthi K, Kawamata H, Rajendran B, Boyer PJ, et al. 2007. Overexpression of CCS in G93A-SOD1 mice leads to accelerated neurological deficits with severe mitochondrial pathology. *Proc. Natl. Acad. Sci. USA* 104:6072–77
136. Kawamata H, Manfredi G. 2008. Different regulation of wild-type and mutant Cu,Zn superoxide dismutase localization in mammalian mitochondria. *Hum. Mol. Genet.* 17:3303–17
137. Reddehase S, Grumbt B, Neupert W, Hell K. 2009. The disulfide relay system of mitochondria is required for the biogenesis of mitochondrial Ccs1 and Sod1. *J. Mol. Biol.* 385:331–38
138. Zhang R, Snyder GH. 1991. Factors governing selective formation of specific disulfides in synthetic variants of  $\alpha$ -conotoxin. *Biochemistry* 30:11343–48
139. Liu H, Zhu H, Eggers DK, Nersissian AM, Faull KF, et al. 2000. Copper<sup>2+</sup> binding to the surface residue cysteine 111 of His46Arg human copper-zinc superoxide dismutase, a familial amyotrophic lateral sclerosis mutant. *Biochemistry* 39:8125–32
140. Peter C, Laliberte J, Beaudoin J, Labbe S. 2008. Copper distributed by Atx1 is available to copper amine oxidase 1 in *Schizosaccharomyces pombe*. *Eukaryot. Cell* 7:1781–94
141. Puig S, Penarrubia L. 2009. Placing metal micronutrients in context: transport and distribution in plants. *Curr. Opin. Plant Biol.* 12:299–306
142. Puig S, Mira H, Dorcey E, Sancenon V, Andres-Colas N, et al. 2007. Higher plants possess two different types of ATX1-like copper chaperones. *Biochem. Biophys. Res. Commun.* 354:385–90
143. Andres-Colas N, Sancenon V, Rodriguez-Navarro S, Mayo S, Thiele DJ, et al. 2006. The *Arabidopsis* heavy metal P-type ATPase HMA5 interacts with metallochaperones and functions in copper detoxification of roots. *Plant J.* 45:225–36
144. Cobine P, Wickramasinghe WA, Harrison MD, Weber T, Solioz M, Dameron CT. 1999. The *Enterococcus hirae* copper chaperone CopZ delivers copper(I) to the CopY repressor. *FEBS Lett.* 445:27–30
145. Cobine PA, George GN, Jones CE, Wickramasinghe WA, Solioz M, Dameron CT. 2002. Copper transfer from the Cu(I) chaperone, CopZ, to the repressor, Zn(II)CopY: metal coordination environments and protein interactions. *Biochemistry* 41:5822–29
146. Itoh S, Kim HW, Nakagawa O, Ozumi K, Lessner SM, et al. 2008. Novel role of antioxidant-1 (Atox1) as a copper dependent transcription factor involved in cell proliferation. *J. Biol. Chem.* 283:9157–67
147. Angeletti B, Waldron KJ, Freeman KB, Bawagan H, Hussain I, et al. 2005. BACE1 cytoplasmic domain interacts with the copper chaperone for superoxide dismutase-1 and binds copper. *J. Biol. Chem.* 280:17930–37



# Contents

## Preface

The Power of One

*James E. Rothman* .....v

## Prefatory Article

Frontispiece

*Aaron Klug* .....xiv

From Virus Structure to Chromatin: X-ray Diffraction  
to Three-Dimensional Electron Microscopy

*Aaron Klug* .....1

## Recent Advances in Biochemistry

Genomic Screening with RNAi: Results and Challenges

*Stephanie Mohr, Chris Bakal, and Norbert Perrimon* .....37

Nanomaterials Based on DNA

*Nadrian C. Seeman* .....65

Eukaryotic Chromosome DNA Replication: Where, When, and How?

*Hisao Masai, Seiji Matsumoto, Zhiying You, Naoko Yoshizawa-Sugata,  
and Masako Oda* .....89

Regulators of the Cohesin Network

*Bo Xiong and Jennifer L. Gerton* .....131

Reversal of Histone Methylation: Biochemical and Molecular

Mechanisms of Histone Demethylases

*Nima Mosammaparast and Yang Shi* .....155

The Mechanism of Double-Strand DNA Break Repair by the  
Nonhomologous DNA End-Joining Pathway

*Michael R. Lieber* .....181

The Discovery of Zinc Fingers and Their Applications in Gene  
Regulation and Genome Manipulation

*Aaron Klug* .....213

Origins of Specificity in Protein-DNA Recognition <i>Remo Robs, Xiangshu Jin, Sean M. West, Robit Joshi, Barry Honig, and Richard S. Mann</i> .....	233
Transcript Elongation by RNA Polymerase II <i>Luke A. Selth, Stefan Sigurdsson, and Jesper Q. Svejstrup</i> .....	271
Biochemical Principles of Small RNA Pathways <i>Qinghua Liu and Zain Paroo</i> .....	295
Functions and Regulation of RNA Editing by ADAR Deaminases <i>Kazuko Nishikura</i> .....	321
Regulation of mRNA Translation and Stability by microRNAs <i>Marc Robert Fabian, Nabum Sonenberg, and Witold Filipowicz</i> .....	351
Structure and Dynamics of a Processive Brownian Motor: The Translating Ribosome <i>Joachim Frank and Ruben L. Gonzalez, Jr.</i> .....	381
Adding New Chemistries to the Genetic Code <i>Chang C. Liu and Peter G. Schultz</i> .....	413
Bacterial Nitric Oxide Synthases <i>Brian R. Crane, Jawabar Sudhamsu, and Bhumit A. Patel</i> .....	445
Enzyme Promiscuity: A Mechanistic and Evolutionary Perspective <i>Olga Khersonsky and Dan S. Tarwfk</i> .....	471
Hydrogenases from Methanogenic Archaea, Nickel, a Novel Cofactor, and H <sub>2</sub> Storage <i>Rudolf K. Thauer, Anne-Kristin Kaster, Meike Goenrich, Michael Schick, Takeshi Hiromoto, and Seigo Shima</i> .....	507
Copper Metallochaperones <i>Nigel J. Robinson and Dennis R. Winge</i> .....	537
High-Throughput Metabolic Engineering: Advances in Small-Molecule Screening and Selection <i>Jeffrey A. Dietrich, Adrienne E. McKee, and Jay D. Keasling</i> .....	563
Botulinum Neurotoxin: A Marvel of Protein Design <i>Mauricio Montal</i> .....	591
Chemical Approaches to Glycobiology <i>Laura L. Kiessling and Rebecca A. Splain</i> .....	619
Cellulosomes: Highly Efficient Nanomachines Designed to Deconstruct Plant Cell Wall Complex Carbohydrates <i>Carlos M.G.A. Fontes and Harry J. Gilbert</i> .....	655



Somatic Mitochondrial DNA Mutations in Mammalian Aging <i>Nils-Göran Larsson</i> .....	683
Physical Mechanisms of Signal Integration by WASP Family Proteins <i>Shae B. Padrick and Michael K. Rosen</i> .....	707
Amphipols, Nanodiscs, and Fluorinated Surfactants: Three Nonconventional Approaches to Studying Membrane Proteins in Aqueous Solutions <i>Jean-Luc Popot</i> .....	737
Protein Sorting Receptors in the Early Secretory Pathway <i>Julia Dancourt and Charles Barlowe</i> .....	777
Virus Entry by Endocytosis <i>Jason Mercer, Mario Schelbaas, and Ari Helenius</i> .....	803

## Indexes

Cumulative Index of Contributing Authors, Volumes 75–79 .....	835
Cumulative Index of Chapter Titles, Volumes 75–79 .....	839

## Errata

An online log of corrections to *Annual Review of Biochemistry* articles may be found at <http://biochem.annualreviews.org>