

Problem Set 8

- For a galvanic cell at 298 K and made using nickel and cadmium electrodes immersed in solutions with a Ni^{2+} concentration of 0.15 M and a Cd^{2+} concentration of 0.50 M respectively:
 - Calculate the voltage for this cell if both solutions were at 1.0 M concentrations.
 - Calculate the voltage for this cell at the given solution concentrations.
 - Write the equation for the oxidation reaction that spontaneously occurs.
 - Write the equation for the reduction reaction that spontaneously occurs.
 - Write the equation for the overall reaction that spontaneously occurs.
 - Write an equation for Q for the overall reaction.
 - Calculate the change in Gibbs Free Energy for the overall reaction.
 - Determine the value of the equilibrium constant for the overall reaction.
 - Draw a diagram of the galvanic cell and show how the electrons flow, along with the reaction that occurs at each electrode.
 - Predict how the masses of each electrode change over time.
 - An Ampere (A) unit is used to measure electrical current; one Ampere represents a current flow of one Coulomb of charge per second. If a current of 10 mA flows for 30 minutes in the galvanic cell above, calculate the change in mass for each electrode.
- Calculate the Nernst potential, in mV, for the bicarbonate ion that had extracellular and intracellular concentrations of 27 mM and 8 mM respectively. For a cell with a resting membrane potential of -70 mV, explain what would happen if a bicarbonate ion channel opened. Would this have an inhibitory or excitatory neural effect?
- A semi-permeable membrane that is permeable to potassium ions, but not to iodide ions, separates a vessel into two compartments. KI solutions of 5.00 mM and 1.00 mM are poured into the two compartments at a temperature of 298K.
 - Calculate the voltage that develops across the membrane.
 - Draw and label a diagram that shows the two compartment concentrations and the sign of the potential gradient across the membrane.
 - Determine what occurs if the 1.00 mM solution is replaced with a 30.00 mM solution. Draw a labeled diagram.